

March, 2010

Hadronic D decays involving even-parity light mesons

Hai-Yang Cheng,^{1,2} Cheng-Wei Chiang^{3,1}

¹ Institute of Physics, Academia Sinica
Taipei, Taiwan 115, Republic of China

² Physics Department, Brookhaven National Laboratory
Upton, New York 11973

³ Department of Physics and Center for Mathematics
and Theoretical Physics, National Central University
Chungli, Taiwan 320, Republic of China

Abstract

We study the hadronic D meson decays into a pseudoscalar meson P and an even-parity meson M , where M represents a scalar meson S , an axial-vector meson A , or a tensor meson T . These decays are first analyzed in the flavor-diagram approach. Fits to the SP modes with S being a non-strange scalar meson show that neither the simple $q\bar{q}$ picture nor the $q^2\bar{q}^2$ scheme is favored by data. Current measurements on the AP decays are insufficient for a meaningful analysis. Some TP data are inconsistent with the others. In certain cases, the W -annihilation diagrams indicated by the data are unexpectedly large. As a comparison, we also compute their decay rates in the factorization approach using form factors extracted from the covariant light-front model. We find that factorization works well for Cabibbo-allowed $D^+ \rightarrow SP, AP$ decays free of the weak annihilation contributions (W -exchange or W -annihilation). For the other SP and AP modes, it is necessary to include weak annihilation contributions to account for the data. However, factorization fails for $D \rightarrow TP$ decays for some unknown reason; the predicted rates are in general too small by at least two orders of magnitude compared to experiment. We also examine the finite width effects of resonances. Some decay modes which are kinematically forbidden become physically allowed due to the finite width of the resonance. We show that the branching fraction of $D^+ \rightarrow \sigma\pi^+$ extracted from three-body decays is enhanced by a factor of 2, whereas $\mathcal{B}(D^0 \rightarrow f_2(1270)\bar{K}^0)$ is reduced by a factor of 4 by finite width effects.

I. INTRODUCTION

A plethora of interesting but puzzling phenomena regarding strong interactions have been revealed in the study of charmed meson decays. The magnitudes and phases of CKM factors associated with the dominant decay amplitudes in such processes are well-determined. Moreover, these decays do not receive significant corrections from penguin-type loop diagrams. Such salient features enable one to readily extract the magnitudes and relative strong phases of various flavor diagrams. More specifically, there are four types of flavor diagrams that dominate the charmed meson decays. They are the color-allowed tree amplitude, the color-suppressed tree amplitude, the W -exchange diagram, and the W -annihilation diagram. Previous studies [1–8] have shown the importance of the W -exchange and W -annihilation diagrams, presumably due to significant final-state rescattering effects, in the two-body decays of charmed mesons to two pseudoscalar (PP) or one vector and one pseudoscalar mesons (VP). Moreover, these amplitudes are seen to have nontrivial relative strong phases, also a result of sizable final-state interactions (FSIs). Among various sources of FSIs, the most important one is arguably the contribution from intermediate resonance states near the D meson masses.

In this paper, we set to study the hadronic decays of charmed mesons into a pseudoscalar meson P and an even-parity meson M . Here M can be a scalar meson, denoted by S , an axial-vector meson, denoted by A , or a tensor meson, denoted by T . The $D \rightarrow SP$ decays have been studied previously in Refs. [9–15]; the $D \rightarrow AP$ decays in Refs. [16–24]; and the $D \rightarrow TP$ decays in Refs. [25–27]. In these decays, the flavor diagram of each topology has two possibilities: one with the spectator quark in the charmed meson going to the pseudoscalar meson in the final state; and the other with the spectator quark ending up in the even-parity meson M . We thus need two copies of each topological diagram to describe the decay processes. Many of these decays have been observed in recent years through dedicated experiments and powerful Dalitz plot analysis of multi-body decays. An extraction of the sizes and relative strong phases of these amplitudes therefore becomes possible.

One purpose of studying these decays is to check our understanding in the structures and properties of light even-parity mesons. Another goal is to learn the FSI pattern in view of the rich resonance spectrum around the D meson mass range.

This paper is organized as follows. In Section II, we review current experimental status of the measurements of multi-body charmed meson decays that are relevant to our analysis. We provide the information of flavor SU(3) classification, decay constants, and form factors for the light S , A , and T mesons in Section III. Section IV presents the so-called flavor-diagram approach to the decays. Each decay mode is decomposed in terms of quark diagrams characterized by their flavor topologies. Current experimental data are used to infer the magnitude and strong phase associated with each of the amplitudes as best as we can. Under the factorization assumption, we compute the rate of each decay mode in Section V. We also examine the finite width effects for certain decay modes in Section VI. Some discussions on the form factor model used in this work, theoretical uncertainties, comparison with other approaches are presented in Section VII. A summary of our findings is given in Section VIII.

II. EXPERIMENTAL STATUS

It is known that three- and four-body decays of heavy mesons provide a rich laboratory for studying the intermediate state resonances. The Dalitz plot analysis is a very useful technique for this purpose. We are interested in $D \rightarrow MP$ decays extracted from the three-body decays of charmed mesons. Many results are available from ARGUS, Femilab, CLEO, FOCUS and BaBar. The results of various experiments are summarized in Tables I-III where the products of $\mathcal{B}(D \rightarrow MP)$ and $\mathcal{B}(M \rightarrow P_1 P_2)$ are listed. To extract the branching fraction for $D \rightarrow MP$, we apply the narrow width approximation

$$\Gamma(D \rightarrow MP \rightarrow P_1 P_2 P) = \Gamma(D \rightarrow MP) \mathcal{B}(M \rightarrow P_1 P_2) . \quad (1)$$

(Finite width effects in certain decays will be discussed in Section VI.) For the branching fractions of two-body decays of even-parity mesons, we shall use

$$\begin{aligned} \mathcal{B}(f_0(980) \rightarrow \pi^+ \pi^-) &= 0.35 \pm 0.08 , & \mathcal{B}(a_0^+(980) \rightarrow \eta \pi) &= 0.845 \pm 0.017 , \\ \mathcal{B}(K_0^{*0}(1430) \rightarrow K^+ \pi^-) &= \frac{2}{3}(0.93 \pm 0.10) , & \mathcal{B}(K_0^{*+}(1430) \rightarrow K^+ \pi^0) &= \frac{1}{3}(0.93 \pm 0.10) , \\ \mathcal{B}(f_2(1270) \rightarrow \pi \pi) &= (84.8_{-1.2}^{+2.4})\% , & \mathcal{B}(f_2(1270) \rightarrow K \bar{K}) &= (4.6 \pm 0.4)\% , \\ \mathcal{B}(a_2(1320) \rightarrow K \bar{K}) &= (4.9 \pm 0.8)\% , & \mathcal{B}(K_2^*(1430) \rightarrow K \pi) &= (49.9 \pm 1.2)\% , \end{aligned} \quad (2)$$

and $\mathcal{B}(f_0(1500) \rightarrow \pi^+ \pi^-) = \frac{2}{3}(34.9 \pm 2.3)\%$, where we have applied the value of $\Gamma(f_0(980) \rightarrow \pi \pi) / [\Gamma(f_0(980) \rightarrow \pi \pi) + \Gamma(f_0(980) \rightarrow K \bar{K})] = 0.52 \pm 0.12$ obtained by BaBar [28] and the Particle Data Group (PDG) average, $\Gamma(a_0(980) \rightarrow K \bar{K}) / \Gamma(a_0(980) \rightarrow \pi \eta) = 0.183 \pm 0.024$ [29]. To obtain the branching fraction of $f_0(980) \rightarrow \pi \pi$, we have assumed that $\pi \pi$ and $K \bar{K}$ are the dominant decay modes of $f_0(980)$.

Several remarks are in order:

1. There are two measurements of $D_s^+ \rightarrow f_0(980) \pi^+$ from $D_s^+ \rightarrow \pi^+ \pi^+ \pi^-$ by E687 [30] and E791 [31] with the results

$$\mathcal{B}(D_s^+ \rightarrow f_0(980) \pi^+) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = \begin{cases} (6.3 \pm 0.8) \times 10^{-3} & \text{E791,} \\ (1.2 \pm 0.2) \times 10^{-2} & \text{E687.} \end{cases} \quad (3)$$

These two data are not used by PDG for the average. The most recent Dalitz plot analysis of $D_s^+ \rightarrow \pi^+ \pi^+ \pi^-$ by BaBar yields $\mathcal{B}(D_s^+ \rightarrow (\pi^+ \pi^-)_{S\text{-wave}} \pi^+) = (0.92 \pm 0.07) \times 10^{-2}$ [32]. Since the S -wave is the sum over $f_0(980)$, $f_0(1370)$ and $f_0(1500)$, it is clear that the E687 result is too large. Hence, we will only quote the E791 result in Table I for $D_s^+ \rightarrow f_0(980) \pi^+$ with $f_0(980) \rightarrow \pi^+ \pi^-$. For an early theoretical study, see [33].

2. Many of the 3-body decays listed in Table I involve the decays $f_0(980) \rightarrow K^+ K^-$ and $a_0(980) \rightarrow K^+ K^-$. Since the central values of the $f_0(980)$ and $a_0(980)$ masses are below the threshold for decay into a pair of charged kaons, the narrow width approximation Eq. (1) is no longer applicable because f_0 or a_0 need to be off-shell. That is, the relation

$$\mathcal{B}(D^0 \rightarrow f_0(980) \pi^0; f_0 \rightarrow K^+ K^-) = \mathcal{B}(D^0 \rightarrow f_0(980) \pi^0) \mathcal{B}(f_0 \rightarrow K^+ K^-) , \quad (4)$$

does not hold. From Table I, we see that $\mathcal{B}(D^0 \rightarrow f_0 \pi^0; f_0 \rightarrow \pi^+ \pi^-)$ is smaller than $\mathcal{B}(D^0 \rightarrow f_0 \pi^0; f_0 \rightarrow K^+ K^-)$ by one order of magnitude. Since $\mathcal{B}(f_0 \rightarrow K^+ K^-)$ is smaller

than $\mathcal{B}(f_0 \rightarrow \pi^+\pi^-)$ due to phase space suppression¹, the branching fraction $\mathcal{B}(f_0(980) \rightarrow K^+K^-)$ obtained from $\mathcal{B}(D^0 \rightarrow f_0\pi^0; f_0 \rightarrow K^+K^-)$ under the narrow width approximation will be too large by at least one order of magnitude.

3. The decay $D^+ \rightarrow K^-\pi^+\pi^+$ is dominated by the S -wave ($K^-\pi^+$) component which consists of $K_0^*(800)$ (or κ), $K_0^*(1430)$ and non-resonant contributions. The PDG value is $\mathcal{B}(D^+ \rightarrow (K^-\pi^+)_{S\text{-wave}}\pi^+) = (7.62 \pm 0.25)\%$ [29]. However, PDG does not take the measurements of $D^+ \rightarrow \bar{K}_0^{*0}(800)\pi^+$ and $\bar{K}_0^{*0}(1430)\pi^+$ by E791, E691 and E687 for the average. If we take the E791 results alone [31], we find

$$\mathcal{B}(D^+ \rightarrow \bar{K}_0^{*0}(800)\pi^+) = (6.5 \pm 1.9)\% , \quad \mathcal{B}(D^+ \rightarrow \bar{K}_0^{*0}(1430)\pi^+) = (1.8 \pm 0.3)\% . \quad (5)$$

Hence, the decay $D^+ \rightarrow \bar{K}_0^{*0}(800)\pi^+$ has the largest branching fraction among the two-body $D \rightarrow SP$ decays.

4. Since the axial-vector mesons decay into three pseudoscalar mesons via strong interactions, their resonant substructures are studied in the Dalitz plot analysis of four-body decays. For example, information on the decay rates of $D^0 \rightarrow K^-a_1^+(1260)$ and $K_1^-(1270)\pi^+$ can be extracted from the study of $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$.
5. Because the σ meson is very broad in its width, of $\mathcal{O}(600 - 1000)$ MeV [29], the use of the narrow width approximation is not justified and it becomes necessary to take into account the finite width effect of σ . We will examine the finite width effect for the decay $D \rightarrow \sigma\pi$ in Section VI.

III. PHYSICAL PROPERTIES OF EVEN-PARITY MESONS

A. Scalar mesons

It is known that the underlying structure of scalar mesons is not well established theoretically (for a review, see *e.g.* Refs. [37, 38]). Many scalar mesons with masses lower than 2 GeV have been observed, and they can be classified into two nonets: one nonet with masses below or close to 1 GeV, namely, the isoscalars $f_0(600)$ (or σ), $f_0(980)$, the isodoublet $K_0^*(800)$ (or κ) and the isovector $a_0(980)$; and the other nonet with masses above 1 GeV, namely, $f_0(1370)$, $a_0(1450)$, $K_0^*(1430)$ and $f_0(1500)/f_0(1710)$ ². If the scalar meson states below or near 1 GeV are identified as a conventional low-lying $0^+ q\bar{q}$ nonet, then the nonet states above 1 GeV could be excited $q\bar{q}$ states.

In the naive quark model, the flavor wave functions of the light scalars read

$$\sigma = \frac{1}{\sqrt{2}}(u\bar{u} + d\bar{d}) , \quad f_0 = s\bar{s} ,$$

¹ The ratio $\mathcal{B}(f_0 \rightarrow K^+K^-)/\mathcal{B}(f_0 \rightarrow \pi^+\pi^-)$ was measured to be 0.69 ± 0.32 by BaBar [28] and $0.25^{+0.17}_{-0.11}$ by BES [34].

² Since not all three isosinglet scalars $f_0(1710)$, $f_0(1500)$, $f_0(1370)$ can be accommodated in the $q\bar{q}$ nonet picture, it is widely believed that one of them should be primarily a scalar glueball.

TABLE I: Experimental branching fractions of various $D \rightarrow SP$ decays. For simplicity and convenience, we have dropped the mass identification for $f_0(980)$, $a_0(980)$ and $K_0^*(1430)$. Data are taken from Ref. [29] unless specified otherwise.

$\mathcal{B}(D \rightarrow SP) \times \mathcal{B}(S \rightarrow P_1 P_2)$	$\mathcal{B}(D \rightarrow SP)$
$\mathcal{B}(D^+ \rightarrow f_0 \pi^+) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (1.55 \pm 0.33) \times 10^{-4}$	$\mathcal{B}(D^+ \rightarrow f_0 \pi^+) = (4.5 \pm 1.4) \times 10^{-4}$
$\mathcal{B}(D^+ \rightarrow f_0(1370) \pi^+) \mathcal{B}(f_0(1370) \rightarrow \pi^+ \pi^-) = (8 \pm 4) \times 10^{-5}$	
$\mathcal{B}(D^+ \rightarrow f_0(1500) \pi^+) \mathcal{B}(f_0(1500) \rightarrow \pi^+ \pi^-) = (1.1 \pm 0.4) \times 10^{-4}$	$\mathcal{B}(D^+ \rightarrow f_0(1500) \pi^+) = (4.7 \pm 1.7) \times 10^{-4}$
$\mathcal{B}(D^+ \rightarrow f_0(1710) \pi^+) \mathcal{B}(f_0(1710) \rightarrow \pi^+ \pi^-) < 5 \times 10^{-5}$	
$\mathcal{B}(D^+ \rightarrow f_0 K^+) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (5.6 \pm 3.4) \times 10^{-5}$	$\mathcal{B}(D^+ \rightarrow f_0 K^+) = (1.6 \pm 1.0) \times 10^{-4}$
$\mathcal{B}(D^+ \rightarrow \sigma \pi^+) \mathcal{B}(\sigma \rightarrow \pi^+ \pi^-) = (1.37 \pm 0.12) \times 10^{-3}$	$\mathcal{B}(D^+ \rightarrow \sigma \pi^+) = (2.1 \pm 0.2) \times 10^{-3}$
$\mathcal{B}(D^+ \rightarrow \bar{\kappa}^0 K^+) \mathcal{B}(\bar{\kappa}^0 \rightarrow K^- \pi^+) = (6.8_{-2.1}^{+3.5}) \times 10^{-4}$	$\mathcal{B}(D^+ \rightarrow \bar{\kappa}^0 K^+) = (1.0_{-0.3}^{+0.5}) \times 10^{-3}$
$\mathcal{B}(D^+ \rightarrow \bar{K}_0^{*0} K^+; \bar{K}_0^{*0} \rightarrow K^- \pi^+) = (1.83 \pm 0.35) \times 10^{-3}$	
$\mathcal{B}(D^+ \rightarrow (K^- \pi^+)_{S\text{-wave}} \pi^+) = (7.62 \pm 0.25)\%$	
$\mathcal{B}(D^0 \rightarrow f_0 \bar{K}^0) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (2.76_{-0.44}^{+0.60}) \times 10^{-3}$	$\mathcal{B}(D^0 \rightarrow f_0 \bar{K}^0) = (8.0_{-2.2}^{+2.5}) \times 10^{-3}$
$\mathcal{B}(D^0 \rightarrow f_0(1370) \bar{K}^0) \mathcal{B}(f_0(1370) \rightarrow \pi^+ \pi^-) = (5.0 \pm 1.2) \times 10^{-3}$	
$\mathcal{B}(D^0 \rightarrow f_0 \pi^0) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (3.6 \pm 0.8) \times 10^{-5}$	$\mathcal{B}(D^0 \rightarrow f_0 \pi^0) = (1.0 \pm 0.3) \times 10^{-4}$
$\mathcal{B}(D^0 \rightarrow f_0 \pi^0; f_0 \rightarrow K^+ K^-) = (3.5 \pm 0.6) \times 10^{-4}$	
$\mathcal{B}(D^0 \rightarrow f_0(1370) \pi^0) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (5.3 \pm 2.1) \times 10^{-5}$	
$\mathcal{B}(D^0 \rightarrow f_0(1500) \pi^0) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (5.6 \pm 1.6) \times 10^{-5}$	$\mathcal{B}(D^0 \rightarrow f_0(1500) \pi^0) = (2.4 \pm 0.7) \times 10^{-4}$
$\mathcal{B}(D^0 \rightarrow f_0(1710) \pi^0) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (4.5 \pm 1.5) \times 10^{-5}$	
$\mathcal{B}(D^0 \rightarrow f_0 \bar{K}^0; f_0 \rightarrow K^+ K^-) < 2.0 \times 10^{-4}$	
$\mathcal{B}(D^0 \rightarrow f_0(1370) \bar{K}^0) \mathcal{B}(f_0 \rightarrow K^+ K^-) = (3.6 \pm 2.2) \times 10^{-4}$	
$\mathcal{B}(D^0 \rightarrow a_0^+ K^-; a_0^+ \rightarrow K^+ \bar{K}^0) = (1.24 \pm 0.36) \times 10^{-3}$	
$\mathcal{B}(D^0 \rightarrow a_0^+ K^-; a_0^+ \rightarrow K^+ \bar{K}^0) = (1.5 \pm 0.1) \times 10^{-3}^a$	
$\mathcal{B}(D^0 \rightarrow a_0^+ (1450) K^-; a_0^+ \rightarrow K^+ \bar{K}^0) = (2.1 \pm 0.1) \times 10^{-3}^a$	
$\mathcal{B}(D^0 \rightarrow a_0^- K^+; a_0^- \rightarrow K^- \bar{K}^0) < 2.4 \times 10^{-4}$	
$\mathcal{B}(D^0 \rightarrow a_0^0 \bar{K}^0; a_0^0 \rightarrow K^+ K^-) = (6.2 \pm 0.8) \times 10^{-3}$	
$\mathcal{B}(D^0 \rightarrow a_0^0 \bar{K}^0; a_0^0 \rightarrow K^+ K^-) = (5.3 \pm 0.4) \times 10^{-3}^a$	
$\mathcal{B}(D^0 \rightarrow a_0^0 \bar{K}^0) \mathcal{B}(a_0^0 \rightarrow \eta \pi^0) = (1.34 \pm 0.42) \times 10^{-2}$	$\mathcal{B}(D^0 \rightarrow a_0^0 \bar{K}^0) = (1.6 \pm 0.5)\%$
$\mathcal{B}(D^0 \rightarrow \sigma \pi^0) \mathcal{B}(\sigma \rightarrow \pi^+ \pi^-) = (1.18 \pm 0.21) \times 10^{-4}$	$\mathcal{B}(D^0 \rightarrow \sigma \pi^0) = (1.8 \pm 0.3) \times 10^{-4}$
$\mathcal{B}(D^0 \rightarrow K_0^{*-} \pi^+) \mathcal{B}(K_0^{*-} \rightarrow \bar{K}^0 \pi^-) = (4.90_{-0.64}^{+0.80}) \times 10^{-3}$	$\mathcal{B}(D^0 \rightarrow K_0^{*-} \pi^+) = (7.9_{-1.3}^{+1.5}) \times 10^{-3}$
$\mathcal{B}(D^0 \rightarrow K_0^{*-} \pi^+) \mathcal{B}(K_0^{*-} \rightarrow K^- \pi^0) = (4.6 \pm 2.1) \times 10^{-3}$	$\mathcal{B}(D^0 \rightarrow K_0^{*-} \pi^+) = (1.5 \pm 0.7)\%$
$\mathcal{B}(D^0 \rightarrow \bar{K}_0^{*0} \pi^0) \mathcal{B}(\bar{K}_0^{*0} \rightarrow K^- \pi^+) = (5.7_{-1.5}^{+5.0}) \times 10^{-3}$	$\mathcal{B}(D^0 \rightarrow \bar{K}_0^{*0} \pi^0) = (9.2_{-2.6}^{+8.1}) \times 10^{-3}$
$\mathcal{B}(D_s^+ \rightarrow (\pi^+ \pi^-)_{S\text{-wave}} \pi^+) = (0.92 \pm 0.07) \times 10^{-2}^b$	
$\mathcal{B}(D_s^+ \rightarrow f_0 \pi^+) \mathcal{B}(f_0 \rightarrow \pi^+ \pi^-) = (6.3 \pm 0.8) \times 10^{-3}^c$	$\mathcal{B}(D_s^+ \rightarrow f_0 \pi^+) = (1.8 \pm 0.5)\%$
$\mathcal{B}(D_s^+ \rightarrow f_0 \pi^+; f_0 \rightarrow K^+ K^-) = (6.0 \pm 2.4) \times 10^{-3}$	
$\mathcal{B}(D_s^+ \rightarrow f_0 \pi^+; f_0 \rightarrow K^+ K^-) = (1.55 \pm 0.13) \times 10^{-2}^d$	
$\mathcal{B}(D_s^+ \rightarrow f_0(1370) \pi^+) \mathcal{B}(f_0 \rightarrow K^+ K^-) = (2.37 \pm 0.35) \times 10^{-2}^d$	
$\mathcal{B}(D_s^+ \rightarrow f_0(1710) \pi^+) \mathcal{B}(f_0 \rightarrow K^+ K^-) = (1.87 \pm 0.29) \times 10^{-2}^d$	
$\mathcal{B}(D_s^+ \rightarrow \bar{K}_0^{*0} K^+) \mathcal{B}(\bar{K}_0^{*0} \rightarrow K^- \pi^+) = (5.1 \pm 2.5) \times 10^{-3}$	$\mathcal{B}(D_s^+ \rightarrow \bar{K}_0^{*0} K^+) = (8.2 \pm 4.1) \times 10^{-3}$
$\mathcal{B}(D_s^+ \rightarrow \bar{K}_0^{*0} K^+) \mathcal{B}(\bar{K}_0^{*0} \rightarrow K^- \pi^+) = (2.15 \pm 0.40) \times 10^{-3}^d$	$\mathcal{B}(D_s^+ \rightarrow \bar{K}_0^{*0} K^+) = (3.5 \pm 0.7) \times 10^{-3}$
$\mathcal{B}(D_s^+ \rightarrow K_0^{*0} \pi^+) \mathcal{B}(K_0^{*0} \rightarrow K^+ \pi^-) = (5 \pm 4) \times 10^{-4}$	$\mathcal{B}(D_s^+ \rightarrow K_0^{*0} \pi^+) = (8.1 \pm 6.5) \times 10^{-4}$

^aFrom BaBar [35].

^bFrom BaBar [32].

^cFrom E791 [31].

^dFrom CLEO [36].

$$\begin{aligned}
 a_0^0 &= \frac{1}{\sqrt{2}}(u\bar{u} - d\bar{d}) , & a_0^+ &= u\bar{d} , & a_0^- &= d\bar{u} , \\
 \kappa^+ &= u\bar{s} , & \kappa^0 &= d\bar{s} , & \bar{\kappa}^0 &= s\bar{d} , & \kappa^- &= s\bar{u} ,
 \end{aligned} \tag{6}$$

where the ideal mixing for f_0 and σ is assumed as $f_0(980)$ is the heaviest one and σ the lightest one in the light scalar nonet. However, this simple picture encounters several serious problems: (i) It is impossible to understand the mass degeneracy between $f_0(980)$ and $a_0(980)$. A related

TABLE II: Experimental branching fractions of $D \rightarrow AP$ decays taken from Ref. [29].

Decay	Experiment	Decay	Experiment
$D^+ \rightarrow \bar{K}_1^0(1270)\pi^+$	$< 7 \times 10^{-3}$	$D^+ \rightarrow \bar{K}^0 a_1^+(1260)$	$(7.0 \pm 1.2)\%$
$D^+ \rightarrow \bar{K}_1^0(1400)\pi^+$	$(3.8 \pm 1.3)\%$	$D^0 \rightarrow K^- a_1^+(1260)$	$(7.9 \pm 1.1)\%$
$D^0 \rightarrow K_1^-(1270)\pi^+$	$(1.14 \pm 0.32)\%$	$D^0 \rightarrow \bar{K}^0 a_1^0(1260)$	$< 1.9\%$
$D^0 \rightarrow K_1^-(1400)\pi^+$	$< 1.2\%$	$D^0 \rightarrow \pi^- a_1^+(1260)$	$(8.98 \pm 0.62) \times 10^{-3}$
$D^0 \rightarrow \bar{K}_1^0(1400)\pi^0$	$< 3.7\%$		
$\mathcal{B}(D^0 \rightarrow K_1^\pm(1270)K^\mp)\mathcal{B}(K_1^\pm(1270) \rightarrow K^\pm\pi^+\pi^-) = (8.1 \pm 1.8) \times 10^{-4}$			
$\mathcal{B}(D^0 \rightarrow K_1^\pm(1400)K^\mp; K_1^\pm(1400) \rightarrow K^\pm\pi^+\pi^-) = (5.4 \pm 1.2) \times 10^{-4}$			

 TABLE III: Experimental branching fractions of various $D \rightarrow TP$ decays. For simplicity and convenience, we have dropped the mass identification for $f_2(1270)$, $a_2(1320)$ and $K_2^*(1430)$. Data are taken from Ref. [29] unless specified otherwise.

$\mathcal{B}(D \rightarrow TP) \times \mathcal{B}(T \rightarrow P_1 P_2)$	$\mathcal{B}(D \rightarrow TP)$
$\mathcal{B}(D^+ \rightarrow f_2\pi^+)\mathcal{B}(f_2 \rightarrow \pi^+\pi^-) = (5.0 \pm 0.9) \times 10^{-4}$	$\mathcal{B}(D^+ \rightarrow f_2\pi^+) = (8.8 \pm 1.6) \times 10^{-4}$
$\mathcal{B}(D^+ \rightarrow \bar{K}_2^{*0}\pi^+)\mathcal{B}(\bar{K}_2^{*0} \rightarrow K^-\pi^+) = (2.1 \pm 0.4) \times 10^{-4}$	$\mathcal{B}(D^+ \rightarrow \bar{K}_2^{*0}\pi^+) = (6.3 \pm 1.2) \times 10^{-4}$
$\mathcal{B}(D^+ \rightarrow K_2^{*0}\pi^+)\mathcal{B}(K_2^{*0} \rightarrow K^+\pi^-) = (5.0 \pm 3.4) \times 10^{-5}$	$\mathcal{B}(D^+ \rightarrow K_2^{*0}\pi^+) = (1.5 \pm 1.0) \times 10^{-4}$
$\mathcal{B}(D^+ \rightarrow \bar{K}_2^{*0}K^+; \bar{K}_2^{*0} \rightarrow K^-\pi^+) = (1.7_{-0.8}^{+1.2}) \times 10^{-4}$	
	$\mathcal{B}(D^+ \rightarrow a_2^+\bar{K}^0) < 3.0 \times 10^{-3}$
$\mathcal{B}(D^0 \rightarrow f_2\pi^0)\mathcal{B}(f_2 \rightarrow \pi^+\pi^-) = (1.91 \pm 0.20) \times 10^{-4}$	$\mathcal{B}(D^0 \rightarrow f_2\pi^0) = (3.4 \pm 0.4) \times 10^{-4}$
$\mathcal{B}(D^0 \rightarrow f_2\bar{K}^0)\mathcal{B}(f_2 \rightarrow \pi^+\pi^-) = (2.8_{-1.2}^{+2.0}) \times 10^{-4}$	$\mathcal{B}(D^0 \rightarrow f_2\bar{K}^0) = (5.0_{-2.1}^{+3.5}) \times 10^{-4}$
$\mathcal{B}(D^0 \rightarrow K_2^{*-}\pi^+)\mathcal{B}(K_2^{*-} \rightarrow \bar{K}^0\pi^-) = (7.0_{-2.2}^{+4.0}) \times 10^{-4}$	$\mathcal{B}(D^0 \rightarrow K_2^{*-}\pi^+) = (2.1_{-0.7}^{+1.2}) \times 10^{-3}$
	$\mathcal{B}(D^0 \rightarrow a_2^+K^-) < 2 \times 10^{-3}$
$\mathcal{B}(D_s^+ \rightarrow f_2\pi^+)\mathcal{B}(f_2 \rightarrow \pi^+\pi^-) = (1.1 \pm 0.2) \times 10^{-3}{}^a$	$\mathcal{B}(D_s^+ \rightarrow f_2\pi^+) = (1.9 \pm 0.4) \times 10^{-3}$
$\mathcal{B}(D_s^+ \rightarrow K_2^{*0}\pi^+)\mathcal{B}(K_2^{*0} \rightarrow K^+\pi^-) = (5 \pm 4) \times 10^{-4}$	$\mathcal{B}(D_s^+ \rightarrow K_2^{*0}\pi^+) = (1.5 \pm 1.2) \times 10^{-3}$

^aFrom BaBar [32].

question is why a_0 is heavier than κ if it does not contain a strange quark? This is the so-called “inverted spectrum problem.” (ii) The P -wave 0^+ meson has a unit of orbital angular momentum which costs energy around 500 MeV. Hence, it should have a higher mass above rather than below 1 GeV. (iii) It is hard to explain why σ and κ are much broader than $f_0(980)$ and $a_0(980)$. (iv) The $\gamma\gamma$ widths of $a_0(980)$ and $f_0(980)$ are much smaller than naively expected for a $q\bar{q}$ state [39]. (v) The radiative decay $\phi \rightarrow a_0(980)\gamma$, which cannot proceed if $a_0(980)$ is a pure $q\bar{q}$ state, can be nicely described in the kaon loop mechanism [40]. This suggests a considerable admixture of the $K\bar{K}$ component.

It turns out that these difficulties can be readily resolved in the tetraquark scenario where the four-quark flavor wave functions of light scalar mesons are symbolically given by [41]

$$\sigma = u\bar{u}d\bar{d}, \quad f_0 = s\bar{s}(u\bar{u} + d\bar{d})/\sqrt{2},$$

$$\begin{aligned}
a_0^0 &= \frac{1}{\sqrt{2}}(u\bar{u} - d\bar{d})s\bar{s} , & a_0^+ &= u\bar{d}s\bar{s} , & a_0^- &= d\bar{u}s\bar{s} , \\
\kappa^+ &= u\bar{s}d\bar{d} , & \kappa^0 &= d\bar{s}u\bar{u} , & \bar{\kappa}^0 &= s\bar{d}u\bar{u} , & \kappa^- &= s\bar{u}d\bar{d} .
\end{aligned} \tag{7}$$

The four quarks $q^2\bar{q}^2$ can form an S -wave (not P -wave!) 0^+ meson without introducing a unit of orbital angular momentum. Moreover, color and spin dependent interactions favor a flavor nonet configuration with attraction between the qq and $\bar{q}\bar{q}$ pairs. Therefore, the 0^+ $q^2\bar{q}^2$ nonet has a mass near or below 1 GeV. This four-quark description explains naturally the inverted mass spectrum of the light nonet, especially the mass degeneracy of $f_0(980)$ and $a_0(980)$. The fall-apart strong decays $\sigma \rightarrow \pi\pi$, $\kappa \rightarrow K\pi$ and $f_0, a_0 \rightarrow K\bar{K}$ are OZI super-allowed without the need of any gluon exchange. This explains the broad widths of σ and κ , while $f_0(980)$ and $a_0(980)$ are narrow because of the suppressed phase space for their decays to the kaon pairs. The decays of $f_0(980)$ and $a_0(980)$ are dominated by $f_0(980) \rightarrow \pi\pi$ and $a_0(980) \rightarrow \eta\pi$, respectively. Lattice calculations have confirmed that $a_0(1450)$ and $K_0^*(1430)$ are $q\bar{q}$ mesons, and suggested that σ and κ are tetraquark mesonia [42, 43]. Since exotic 4-quark states have not been seen experimentally, this may imply the structure of diquark-antidiquark bound states for the light scalar mesons (for a review, see Ref. [44]).

In the 2-quark picture with ideal mixing, $f_0(980)$ is purely an $s\bar{s}$ state. This is supported by the data of $D_s^+ \rightarrow f_0\pi^+$ and $\phi \rightarrow f_0\gamma$, implying the copious $f_0(980)$ production via its $s\bar{s}$ component. However, there also exists some experimental evidence indicating that $f_0(980)$ is not a pure $s\bar{s}$ state. First, the observation of $\Gamma(J/\psi \rightarrow f_0\omega) \approx \frac{1}{2}\Gamma(J/\psi \rightarrow f_0\phi)$ [29] clearly shows the existence of the non-strange and strange quark contents in $f_0(980)$. Second, the facts that $f_0(980)$ and $a_0(980)$ have similar widths and that the f_0 width is dominated by $\pi\pi$ also suggest the composition of $u\bar{u}$ and $d\bar{d}$ pairs in $f_0(980)$; that is, $f_0(980) \rightarrow \pi\pi$ should not be OZI suppressed relative to $a_0(980) \rightarrow \pi\eta$. Therefore, isoscalars $\sigma(600)$ and $f_0(980)$ should have a mixing

$$|f_0(980)\rangle = |s\bar{s}\rangle \cos \theta + |n\bar{n}\rangle \sin \theta , \quad |\sigma(600)\rangle = -|s\bar{s}\rangle \sin \theta + |n\bar{n}\rangle \cos \theta , \tag{8}$$

with $n\bar{n} \equiv (\bar{u}u + \bar{d}d)/\sqrt{2}$. Experimental implications for the f_0 - σ mixing angle have been discussed in detail in Ref. [13]: the mixing angle lies in the ranges of $25^\circ < \theta < 40^\circ$ and $140^\circ < \theta < 165^\circ$.

Likewise, in the four-quark scenario for light scalar mesons, one can also define a similar f_0 - σ mixing angle

$$|f_0(980)\rangle = |n\bar{n}s\bar{s}\rangle \cos \phi + |u\bar{u}d\bar{d}\rangle \sin \phi , \quad |\sigma(600)\rangle = -|n\bar{n}s\bar{s}\rangle \sin \phi + |u\bar{u}d\bar{d}\rangle \cos \phi . \tag{9}$$

It has been shown that $\phi = 174.6^\circ$ [47].

³ Recently CLEO has measured the semileptonic decay $D_s^+ \rightarrow f_0(980)e^+\nu_e$ with the result $\mathcal{B}(D_s^+ \rightarrow f_0(980)e^+\nu_e)\mathcal{B}(f_0 \rightarrow \pi^+\pi^-) = (0.20 \pm 0.03 \pm 0.01)\%$ [45]. Using the value $\mathcal{B}(f_0 \rightarrow \pi^+\pi^-) = (50_{-9}^{+7})\%$ inferred from the BES measurement [34] and the QCD sum rule prediction $\mathcal{B}(D_s^+ \rightarrow f_0e^+\nu) = \cos^2 \theta \times (0.41)\%$ [46], CLEO then extracted the mixing angle to be $\cos^2 \theta = 0.98_{-0.21}^{+0.02}$. However, this is subject to two major uncertainties. First, the branching fraction of $f_0(980) \rightarrow \pi^+\pi^-$ has not been measured directly. For $\mathcal{B}(f_0(980) \rightarrow \pi^+\pi^-) \approx 0.35$ as used in this work, $\mathcal{B}(D_s^+ \rightarrow f_0(980)e^+\nu_e)$ and $\cos^2 \theta$ would be enhanced by a factor of 1.4. Second, the theoretical prediction of this semileptonic decay is model dependent as it depends on the form factor of the $D_s^+ \rightarrow f_0^s$ transition with $f_0^s = s\bar{s}$.

In principle, the 2-quark and 4-quark descriptions of the light scalars can be discriminated in the semileptonic charm decays. For example, the ratio

$$R = \frac{\mathcal{B}(D^+ \rightarrow f_0 \ell^+ \nu) + \mathcal{B}(D^+ \rightarrow \sigma \ell^+ \nu)}{\mathcal{B}(D^+ \rightarrow a_0^0 \ell^+ \nu)} \quad (10)$$

is equal to 1 in the 2-quark scenario and 3 in the 4-quark model under the flavor SU(3) symmetry [48]. In reality, the light scalar mesons may have both 2-quark and 4-quark components. Indeed, a real hadron in the QCD language should be described by a set of Fock states each of which has the same quantum number as the hadron. For example,

$$|a^+(980)\rangle = \psi_{u\bar{d}}^{a_0}|u\bar{d}\rangle + \psi_{u\bar{d}g}^{a_0}|u\bar{d}g\rangle + \psi_{u\bar{d}s\bar{s}}^{a_0}|u\bar{d}s\bar{s}\rangle + \dots \quad (11)$$

In the tetraquark model, $\psi_{u\bar{d}s\bar{s}}^{a_0} \gg \psi_{u\bar{d}}^{a_0}$, while it is the other way around in the 2-quark model.

The decay constant of the scalar meson is defined as⁴

$$\langle S(p) | \bar{q}_2 \gamma_\mu q_1 | 0 \rangle = f_S p_\mu, \quad \langle S | \bar{q}_2 q_1 | 0 \rangle = m_S \bar{f}_S. \quad (12)$$

The neutral scalar mesons σ , f_0 and a_0^0 cannot be produced via the vector current owing to charge conjugation invariance or conservation of vector current:

$$f_\sigma = f_{f_0} = f_{a_0^0} = 0. \quad (13)$$

Applying the equation of motion to Eq. (12) yields

$$\mu_S f_S = \bar{f}_S, \quad \text{with } \mu_S = \frac{m_S}{m_2(\mu) - m_1(\mu)}, \quad (14)$$

where m_2 and m_1 are the running current quark masses. Therefore, the vector decay constant of the scalar meson f_S vanishes in the SU(3) or isospin limit. The vector decay constants of $K_0^*(1430)$ and the charged $a_0(980)$ are non-vanishing, but they are suppressed due to the small mass difference between the constituent s and u quarks and between d and u quarks, respectively. The scalar decay constants $\bar{f}_S$ have been computed in Ref. [49] within the framework of QCD sum rules. From Eq. (14) we obtain $f_{a_0(980)^\pm} = 1.0 \text{ MeV}$, $f_{a_0(1450)^\pm} = 5.3 \text{ MeV}$, and $f_{K_0^*(1430)} = 35.9 \text{ MeV}$. In short, the vector decay constants of scalar mesons are either zero or small.

Form factors for $D \rightarrow P, S$ transitions are defined by [50]

$$\begin{aligned} \langle P(p') | V_\mu | D(p) \rangle &= \left(P_\mu - \frac{m_D^2 - m_P^2}{q^2} q_\mu \right) F_1^{DP}(q^2) + \frac{m_D^2 - m_P^2}{q^2} q_\mu F_0^{DP}(q^2), \\ \langle S(p') | A_\mu | D(p) \rangle &= -i \left[\left(P_\mu - \frac{m_D^2 - m_S^2}{q^2} q_\mu \right) F_1^{DS}(q^2) + \frac{m_D^2 - m_S^2}{q^2} q_\mu F_0^{DS}(q^2) \right], \end{aligned} \quad (15)$$

where $P_\mu = (p + p')_\mu$, $q_\mu = (p - p')_\mu$. As shown in Ref. [51], a factor of $(-i)$ is needed in the $D \rightarrow S$ transition in order for the $D \rightarrow S$ form factors to be positive. This can also be checked from heavy quark symmetry consideration [51].

Throughout this paper, we use the 3-parameter parametrization

$$F(q^2) = \frac{F(0)}{1 - a(q^2/m_D^2) + b(q^2/m_D^2)^2} \quad (16)$$

for $D \rightarrow M$ transitions. The parameters $F(0)$, a and b for $D \rightarrow S$ transitions calculated in the covariant light-front (CLF) quark model are exhibited in Table IV.

⁴ For pseudoscalar mesons, the decay constant is defined as $\langle P(p) | \bar{q}_2 \gamma_\mu \gamma_5 q_1 | 0 \rangle = -i f_P p_\mu$.

TABLE IV: Parameters in the form factors of $D, D_s \rightarrow f_0(980), K_0^*(1430)$ transitions in the parametrization of Eq. (16), as obtained by fitting to the covariant light-front model [51]. The numbers in parentheses are the form factors at $q^2 = 0$ obtained using the ISGW2 model [52].

F	$F(0)$	a	b	F	$F(0)$	a	b
$F_0^{Df_0q}$	0.49 (0.13)	0.07	-0.03	$F_0^{D_s f_0s}$	0.46 (0.23)	-0.29	0.07
$F_0^{DK_0^*(1430)}$	0.48 (0.08)	-0.11	0.02	$F_0^{D_s K_0^*(1430)}$	0.51 (0.14)	0.07	0.02

B. Axial-vector mesons

In the quark model, two nonets of $J^P = 1^+$ axial-vector mesons are expected as the orbital excitation of the $q\bar{q}$ system. In terms of the spectroscopic notation $^{2S+1}L_J$, there are two types of P -wave axial-vector mesons, namely, 3P_1 and 1P_1 . These two nonets have distinctive C quantum numbers for the corresponding neutral mesons, $C = +$ and $C = -$, respectively. Experimentally, the $J^{PC} = 1^{++}$ nonet consists of $a_1(1260)$, $f_1(1285)$, $f_1(1420)$ and K_{1A} , while the 1^{+-} nonet contains $b_1(1235)$, $h_1(1170)$, $h_1(1380)$ and K_{1B} . The physical mass eigenstates $K_1(1270)$ and $K_1(1400)$ are mixtures of the K_{1A} and K_{1B} states (we follow PDG [29] to denote the 3P_1 and 1P_1 states of K_1 by K_{1A} and K_{1B} , respectively),

$$\begin{aligned} K_1(1270) &= K_{1A} \sin \theta_{K_1} + K_{1B} \cos \theta_{K_1} , \\ K_1(1400) &= K_{1A} \cos \theta_{K_1} - K_{1B} \sin \theta_{K_1} . \end{aligned} \quad (17)$$

Since these states are not charge conjugation eigenstates, consequently, mixing is not prohibited. Indeed, the mixing is governed by the mass difference between the strange and non-strange light quarks. There exist several estimations on the mixing angle θ_{K_1} in the literature. From the early experimental information on masses and the partial rates of $K_1(1270)$ and $K_1(1400)$, Suzuki found two possible solutions, each with a two-fold ambiguity, $|\theta_{K_1}| \approx 33^\circ$ and 57° [53]. A similar constraint $35^\circ \lesssim |\theta_{K_1}| \lesssim 55^\circ$ was obtained in Ref. [54] based solely on two parameters: the mass difference between the a_1 and b_1 mesons and the ratio of the constituent quark masses. An analysis of $\tau \rightarrow K_1(1270)\nu_\tau$ and $K_1(1400)\nu_\tau$ decays also yielded the mixing angle to be $\approx 37^\circ$ or 58° with a two-fold ambiguity [55]. Most of these estimations were obtained by assuming a vanishing $f_{K_{1B}}$. With the help of analytical expressions of $f_{K_{1A,1B}}$ obtained in the CLF quark model [51], two solutions for the $K_1(1270)$ - $K_1(1400)$ mixing angle, 50.8° and -44.8° , have been found in Ref. [56]. However, the second solution has been ruled out by the measurements of $B \rightarrow K_1(1270)\gamma$ and $B \rightarrow K_1(1400)\gamma$ [56]. Therefore, we shall use $\theta_{K_1} = 50.8^\circ$ in the ensuing discussions.

For the decay constants and the form factors of the axial vector mesons, we shall follow Ref. [51] to define them as⁵

$$\langle A(p, \varepsilon) | A_\mu | 0 \rangle = f_A m_A \epsilon_\mu^* ,$$

⁵ The relative signs of the decay constants, form factors and mixing angles of the axial-vector mesons were often very confusing in the literature. As stressed in Ref. [57], the sign of the mixing angle θ_{K_1} is intimately related to the relative sign of the K_{1A} and K_{1B} states. In the CLF quark model [51] and in pQCD [58], the decay constants of K_{1A} and K_{1B} are of opposite signs, while the $D(B) \rightarrow K_{1A}$ and

TABLE V: Parameters in the form factors of $D \rightarrow a_1(1260), b_1(1235), K_{1A}, K_{1B}$ transitions in the parametrization of Eq. (16), as obtained by fitting to the covariant light-front model [51]. The numbers in parentheses are the form factors at $q^2 = 0$ obtained using the ISGW2 model [52]. As noticed in the footnote of this subsection, the form factors for $D \rightarrow {}^3P_1$ and $D \rightarrow {}^1P_1$ transitions are of the same (opposite) signs in the CLF (ISGW) model.

F	$F(0)$	a	b	F	$F(0)$	a	b
$V_0^{Da_1}$	0.31 (−0.60)	0.85	0.49	$V_0^{Db_1}$	0.49 (0.64)	0.89	0.28
$V_0^{DK_{1A}}$	0.34 (−0.37)	1.44	0.15	$V_0^{DK_{1B}}$	0.44 (0.50)	0.80	0.27

$$\begin{aligned}
\langle A(p, \varepsilon) | A_\mu | D(p_D) \rangle &= \frac{2}{m_D - m_A} \epsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu} p_D^\alpha p^\beta A^{DA}(q^2), \\
\langle A(p, \varepsilon) | V_\mu | D(p_D) \rangle &= -i \left\{ (m_D - m_A) \epsilon_\mu^* V_1^{DA}(q^2) - (\epsilon^* \cdot p_D)(p_D + p)_\mu \frac{V_2^{DA}(q^2)}{m_D - m_A} \right. \\
&\quad \left. - 2m_A \frac{\epsilon^* \cdot p_D}{q^2} q_\mu [V_3^{DA}(q^2) - V_0^{DA}(q^2)] \right\}. \tag{18}
\end{aligned}$$

Because of the charge conjugation invariance, the decay constant of the 1P_1 non-strange neutral meson such as $b_1^0(1235)$ must be zero. In the isospin limit, the decay constant of the charged b_1 vanishes due to the fact that b_1 has an even G -parity and that the relevant weak axial-vector current is odd under G transformation. As for the strange axial vector mesons, it is known that the decay constant of the 1P_1 meson vanishes in the SU(3) limit [53].

In the following, we shall take $f_{a_1} = 238 \pm 10$ MeV obtained using the QCD sum rule method [61], similar to the ρ meson, $f_\rho \approx 216$ MeV. This means that the $a_1(1260)$ meson can be regarded as the scalar partner of the ρ meson, as it should be. In the CLF quark model [51], if we increase the constituent d quark mass by an amount of 5 ± 2 MeV relative to the u quark mass, we find $f_{b_1} = 0.6 \pm 0.2$ MeV for the charged b_1 which is very small. Using the experimental results $\mathcal{B}(\tau \rightarrow K_1(1270)\nu_\tau) = (4.7 \pm 1.1) \times 10^{-3}$ and $\Gamma(\tau \rightarrow K_1(1270)\nu_\tau)/[\Gamma(\tau \rightarrow K_1(1270)\nu_\tau) + \Gamma(\tau \rightarrow K_1(1400)\nu_\tau)] = 0.69 \pm 0.15$ [29], we obtain

$$|f_{K_1(1270)}| = 169.5^{+18.8}_{-21.2} \text{ MeV}, \quad |f_{K_1(1400)}| = 139.2^{+41.3}_{-45.6} \text{ MeV}. \tag{19}$$

In the CLF quark model the signs of the decay constants $f_{K_{1A}}$ and $f_{K_{1B}}$ are fixed: $f_{K_{1A}} = -212$ MeV and $f_{K_{1B}} = 12$ MeV [56]. This together with the mixing angle $\theta_{K_1} = 50.8^\circ$ also fixes the signs of f_{K_1} to be

$$f_{K_1(1270)} = -170 \text{ MeV}, \quad f_{K_1(1400)} = -139 \text{ MeV}, \tag{20}$$

where we just consider the central values.

Finally, the $D \rightarrow A$ form factor parameters in the CLF quark model are given in Table V.

$D(B) \rightarrow K_{1B}$ form factors are of the same sign. The mixing angle θ_{K_1} is positive. It is the other way around in the approaches of QCD sum rules [59] and the ISGW model [52, 60]: the decay constants of K_{1A} and K_{1B} have the same sign, while the $D(B) \rightarrow K_{1A}$ and $D(B) \rightarrow K_{1B}$ form factors are opposite in sign. These two conventions are related via a redefinition of the K_{1A} or K_{1B} state, *i.e.*, $K_{1A} \rightarrow -K_{1A}$ or $K_{1B} \rightarrow -K_{1B}$.

C. Tensor mesons

The observed $J^P = 2^+$ tensor mesons $f_2(1270)$, $f_2'(1525)$, $a_2(1320)$ and $K_2^*(1430)$ form an $SU(3)$ 1^3P_2 nonet. The $q\bar{q}$ content for isodoublet and isovector tensor resonances are obvious. Just as the η - η' mixing in the pseudoscalar case, the isoscalar tensor states $f_2(1270)$ and $f_2'(1525)$ also have a mixing, and their wave functions are defined by

$$\begin{aligned} f_2(1270) &= \frac{1}{\sqrt{2}}(f_2^u + f_2^d) \cos \theta_{f_2} + f_2^s \sin \theta_{f_2} , \\ f_2'(1525) &= \frac{1}{\sqrt{2}}(f_2^u + f_2^d) \sin \theta_{f_2} - f_2^s \cos \theta_{f_2} , \end{aligned} \quad (21)$$

with $f_2^q \equiv q\bar{q}$. Since $\pi\pi$ is the dominant decay mode of $f_2(1270)$ whereas $f_2'(1525)$ decays predominantly into $K\bar{K}$ (see Ref. [29]), it is obvious that this mixing angle should be small. More precisely, it is found that $\theta_{f_2} = 7.8^\circ$ [62] and $(9 \pm 1)^\circ$ [29]. Therefore, $f_2(1270)$ is primarily an $(u\bar{u} + d\bar{d})/\sqrt{2}$ state, while $f_2'(1525)$ is dominantly $s\bar{s}$.

The polarization tensor $\varepsilon_{\mu\nu}$ of a 3P_2 tensor meson with $J^{PC} = 2^{++}$ satisfies the relations

$$\varepsilon_{\mu\nu} = \varepsilon_{\nu\mu} , \quad \varepsilon^\mu{}_\mu = 0 , \quad p_\mu \varepsilon^{\mu\nu} = p_\nu \varepsilon^{\mu\nu} = 0 , \quad (22)$$

where p^μ is the momentum of the tensor meson. Therefore,

$$\langle 0 | (V - A)_\mu | T(\varepsilon, p) \rangle = a \varepsilon_{\mu\nu} p^\nu + b \varepsilon^\nu{}_\nu p_\mu = 0 , \quad (23)$$

and hence the decay constant of the tensor meson vanishes identically; that is, the tensor meson cannot be produced from the $V - A$ current.

The general expression for the $D \rightarrow T$ transition has the form [60]

$$\begin{aligned} \langle T(\varepsilon, p_T) | (V - A)_\mu | D(p_D) \rangle &= i h(q^2) \varepsilon_{\mu\nu\rho\sigma} \varepsilon^{*\nu\alpha} p_{D\alpha} (p_D + p_T)^\rho (p_D - p_T)^\sigma + k(q^2) \varepsilon_{\mu\nu}^* p_D^\nu \\ &+ b_+(q^2) \varepsilon_{\alpha\beta}^* p_D^\alpha p_D^\beta (p_D + p_T)_\mu + b_-(q^2) \varepsilon_{\alpha\beta}^* p_D^\alpha p_D^\beta (p_D - p_T)_\mu . \end{aligned} \quad (24)$$

The form factors h , k , b_+ and b_- have been calculated in the ISGW quark model [60] and its improved version, the ISGW2 model [52]. They are also computed in the CLF quark model [51] and listed in Table VI.

The decay amplitude of $D \rightarrow TP$ always has the generic expression

$$A(D \rightarrow TP) = \varepsilon_{\mu\nu}^* p_D^\mu p_D^\nu M(D \rightarrow TP) . \quad (25)$$

The decay rate is given by

$$\Gamma(D \rightarrow TP) = \frac{p_c^5}{12\pi m_T^2} \left(\frac{m_D}{m_T} \right)^2 |M(D \rightarrow TP)|^2 , \quad (26)$$

where p_c is the magnitude of the 3-momentum of either final-state meson in the rest frame of the charmed meson.

IV. DIAGRAMMATIC APPROACH

It has been established sometime ago that a least model-dependent analysis of heavy meson decays can be carried out in the so-called topological diagram approach. In this diagrammatic

TABLE VI: Parameters in the form factors of $D \rightarrow a_2(132), K_2^*(1430)$ transitions in the parametrization of Eq. (16), as obtained by fitting to the covariant light-front model [51]. The form factor k is dimensionless, while k , b_+ and b_- are in units of GeV^{-2} . The numbers in parentheses are the form factors at $q^2 = 0$ obtained using the ISGW2 model [52].

F	$F(0)$	a	b	F	$F(0)$	a	b
h^{Da_2}	0.188 (0.203)	1.21	1.09	k^{Da_2}	0.340 (0.613)	-0.07	0.12
$b_+^{Da_2}$	-0.084 (-0.052)	0.97	0.58	$b_-^{Da_2}$	0.120 (0.064)	1.15	0.66
$h^{Df_{2q}}$	0.17 (0.20)	1.28	0.90	$k^{Df_{2q}}$	0.27 (0.61)	-0.21	0.12
$b_+^{Df_{2q}}$	-0.08 (-0.05)	1.02	0.51	$b_-^{Df_{2q}}$	0.10 (0.06)	1.14	0.49
$h^{DK_2^*}$	0.192 (0.14)	1.17	0.99	$k^{DK_2^*}$	0.368 (0.71)	-0.04	0.11
$b_+^{DK_2^*}$	-0.096 (-0.060)	1.05	0.58	$b_-^{DK_2^*}$	0.137 (0.069)	1.17	0.69
$h^{Dsf_{2s}}$	0.15 (0.21)	1.04	0.79	$k^{Dsf_{2s}}$	0.59 (1.15)	0.22	0.09
$b_+^{Dsf_{2s}}$	-0.09 (-0.09)	0.95	0.54	$b_-^{Dsf_{2s}}$	0.13 (0.12)	1.05	0.60
$h^{D_s K_2^*}$	0.15 (0.35)	1.11	0.99	$k^{D_s K_2^*}$	0.42 (1.16)	0.08	0.11
$b_+^{D_s K_2^*}$	-0.07 (-0.08)	0.96	0.60	$b_-^{D_s K_2^*}$	0.11 (0.13)	1.10	0.63

scenario, all two-body nonleptonic weak decays of heavy mesons can be expressed in terms of six distinct quark diagrams [63–65]: T , the color-allowed external W -emission tree diagram; C , the color-suppressed internal W -emission diagram; E , the W -exchange diagram; A , the W -annihilation diagram; P , the horizontal W -loop diagram; and V , the vertical W -loop diagram. (The one-gluon exchange approximation of the P graph is the so-called “penguin diagram”.) It should be stressed that these diagrams are classified according to the topologies of weak interactions with all strong interaction effects encoded, and hence they are *not* Feynman graphs. All quark graphs used in this approach are topological and meant to have all the strong interactions included, *i.e.*, gluon lines are included implicitly in all possible ways. Therefore, analyses of topological graphs can provide information on FSIs. Various topological amplitudes in two-body hadronic D decays have been extracted from the data in [2, 6, 66–72] after making some reasonable approximations, *e.g.*, flavor $\text{SU}(3)$ symmetry.

The topological amplitudes for $D \rightarrow SP, AP, TP$ decays have been discussed in [13, 22, 27]. There are several new features. First, one generally has two sets of distinct external W -emission and internal W -emission diagrams, depending on whether the emitted particle is an even-parity meson or an odd-parity one. Let us denote the primed amplitudes T' and C' for the case when the emitted meson is an even-parity one. Second, because of the smallness of the decay constants of even-parity mesons except for the 3P_1 axial-vector state, it is expected that $|T'| \ll |T|$ and $|C'| \ll |C|$. This feature can be tested experimentally. Third, since K_0^* and the light scalars σ , κ , f_0 , a_0 fall into two different $\text{SU}(3)$ flavor nonets, in principle one cannot apply $\text{SU}(3)$ symmetry to relate the topological amplitudes in $D^+ \rightarrow f_0 \pi^+$ to, for example, those in $D^+ \rightarrow \overline{K}_0^{*0} \pi^+$.

TABLE VII: Topological amplitudes and branching fractions for various $D \rightarrow SP$ decays. In Scheme I, light scalar mesons σ , κ , $a_0(980)$ and $f_0(980)$ are described by the $q\bar{q}$ states, while K_0^* as excited $q\bar{q}$ states. In Scheme II, light scalars are tetraquark states, while K_0^* are ground-state $q\bar{q}$. The $f_0 - \sigma$ mixing angle θ in the 2-quark model is defined in Eq. (8). The experimental branching fractions for $D^0 \rightarrow K_0^{*-}\pi^+$ and $D_s^+ \rightarrow \bar{K}_0^{*0}K^+$ are taken from Table I after average. For simplicity, we do not consider the $f_0 - \sigma$ mixing in the 4-quark model.

Decay	Amplitude (I)	Amplitude (II)	$\mathcal{B}_{\text{expt}}$
$D^0 \rightarrow f_0\pi^0$	$\frac{1}{2}V_{cd}^*V_{ud}(-C + C' - E - E')\sin\theta$ $+\frac{1}{\sqrt{2}}V_{cs}^*V_{us}C'\cos\theta$	$\frac{1}{2}V_{cd}^*V_{ud}(-C + C' - E - E')$ $+V_{cs}^*V_{us}C'$	$(1.0 \pm 0.3) \times 10^{-4}$
$\rightarrow f_0\bar{K}^0$	$V_{cs}^*V_{ud}[\frac{1}{\sqrt{2}}(C + E)\sin\theta + E'\cos\theta]$	$\frac{1}{\sqrt{2}}V_{cs}^*V_{ud}(C + 2E' + E)$	$(8.0_{-2.2}^{+2.5}) \times 10^{-3}$
$\rightarrow a_0^+K^-$	$V_{cs}^*V_{ud}(T' + E)$	$V_{cs}^*V_{ud}(T' + E)$	
$\rightarrow a_0^0\bar{K}^0$	$V_{cs}^*V_{ud}(C - E)/\sqrt{2}$	$V_{cs}^*V_{ud}(C - E)/\sqrt{2}$	$(1.6 \pm 0.5)\%$
$\rightarrow a_0^-K^+$	$V_{cd}^*V_{us}(T + E')$	$V_{cd}^*V_{us}(T + E')$	
$\rightarrow a_0^+\pi^-$	$V_{cd}^*V_{ud}(T' + E)$	$V_{cd}^*V_{ud}(T' + E)$	
$\rightarrow a_0^-\pi^+$	$V_{cd}^*V_{ud}(T + E')$	$V_{cd}^*V_{ud}(T + E')$	
$\rightarrow \sigma\pi^0$	$\frac{1}{2}V_{cd}^*V_{ud}(-C + C' - E - E')\cos\theta$ $-\frac{1}{\sqrt{2}}V_{cs}^*V_{us}C'\sin\theta$	$\frac{1}{\sqrt{2}}V_{cd}^*V_{ud}(-C + C' - E - E')$	$(1.8 \pm 0.3) \times 10^{-4}$
$D^+ \rightarrow f_0\pi^+$	$\frac{1}{\sqrt{2}}V_{cd}^*V_{ud}(T + C' + A + A')\sin\theta$ $+V_{cs}^*V_{us}C'\cos\theta$	$\frac{1}{\sqrt{2}}V_{cd}^*V_{ud}(T + C' + A + A')$ $+\sqrt{2}V_{cs}^*V_{us}C'$	$(4.5 \pm 1.4) \times 10^{-4}$
$\rightarrow f_0K^+$	$V_{cd}^*V_{us}[\frac{1}{\sqrt{2}}(T + A')\sin\theta + A\cos\theta]$	$\frac{1}{\sqrt{2}}V_{cd}^*V_{us}(T + 2A + A')$	$(1.6 \pm 1.0) \times 10^{-4}$
$\rightarrow a_0^+\bar{K}^0$	$V_{cs}^*V_{ud}(T' + C)$	$V_{cs}^*V_{ud}(T' + C)$	
$\rightarrow a_0^0\pi^+$	$V_{cd}^*V_{ud}(-T - C' - A + A')/\sqrt{2}$	$V_{cd}^*V_{ud}(-T - C' - A + A')/\sqrt{2}$	
$\rightarrow \sigma\pi^+$	$\frac{1}{\sqrt{2}}V_{cd}^*V_{ud}(T + C' + A + A')\cos\theta$ $-V_{cs}^*V_{us}C'\sin\theta$	$V_{cd}^*V_{ud}(T + C' + A + A')$	$(2.1 \pm 0.2) \times 10^{-3}$
$\rightarrow \bar{\kappa}^0K^+$	$V_{cs}^*V_{us}T + V_{cd}^*V_{ud}A$	$V_{cs}^*V_{us}T + V_{cd}^*V_{ud}A$	$(1.0_{-0.3}^{+0.5}) \times 10^{-3}$
$D_s^+ \rightarrow f_0\pi^+$	$V_{cs}^*V_{ud}(T\cos\theta + (A + A')\sin\theta/\sqrt{2})$	$V_{cs}^*V_{ud}(2T + A + A')/\sqrt{2}$	$(1.8 \pm 0.5)\%$
$\rightarrow f_0K^+$	$V_{cs}^*V_{us}[(T + C' + A)\cos\theta + \frac{1}{\sqrt{2}}A'\sin\theta]$ $+\frac{1}{\sqrt{2}}V_{cd}^*V_{ud}C'\sin\theta$	$\frac{1}{\sqrt{2}}V_{cs}^*V_{us}(2T + 2C' + 2A + A')$ $+\frac{1}{\sqrt{2}}V_{cd}^*V_{ud}C'$	
$D^0 \rightarrow K_0^{*-}\pi^+$	$V_{cs}^*V_{ud}(T + E')$	$V_{cs}^*V_{ud}(T + E')$	$(8.2 \pm 1.4) \times 10^{-3}$
$\rightarrow \bar{K}_0^{*0}\pi^0$	$V_{cs}^*V_{ud}(C' - E')/\sqrt{2}$	$V_{cs}^*V_{ud}(C' - E')/\sqrt{2}$	$(9.2_{-2.6}^{+8.1}) \times 10^{-3}$
$D^+ \rightarrow \bar{K}_0^{*0}\pi^+$	$V_{cs}^*V_{ud}(T + C')$	$V_{cs}^*V_{ud}(T + C')$	$(1.8 \pm 0.3)\%^a$
$D_s^+ \rightarrow \bar{K}_0^{*0}K^+$	$V_{cs}^*V_{ud}(C' + A)$	$V_{cs}^*V_{ud}(C' + A)$	$(3.6 \pm 0.7) \times 10^{-3}$
$\rightarrow K_0^{*0}\pi^+$	$V_{cd}^*V_{ud}T + V_{cs}^*V_{us}A$	$V_{cd}^*V_{ud}T + V_{cs}^*V_{us}A$	$(8.1 \pm 6.5) \times 10^{-4}$

^aData from E791 [31]; see also Eq. (5).

A. $D \rightarrow SP$

The topological amplitudes for $D \rightarrow SP$ decays are listed in Table VII for two different schemes. In Scheme I, light scalar mesons $\sigma, \kappa, a_0(980)$ and $f_0(980)$ are described by the ground-state $q\bar{q}$ states, while K_0^* as excited $q\bar{q}$ states. In Scheme II, light scalars are tetraquark states, while K_0^* are ground-state $q\bar{q}$. The expressions of topological amplitudes are the same in both Schemes I and II except for the channels involving f_0 or σ .

Since the decay constant of f_0 and σ vanishes, one can set $T' = C' = 0$. From Table VII we have

$$\frac{\mathcal{B}(D^+ \rightarrow f_0\pi^+)}{\mathcal{B}(D^+ \rightarrow \sigma\pi^+)} = \frac{\mathcal{B}(D^0 \rightarrow f_0\pi^0)}{\mathcal{B}(D^0 \rightarrow \sigma\pi^0)} \simeq \begin{cases} \tan^2\theta & \text{2-quark,} \\ \frac{1}{2} & \text{4-quark.} \end{cases} \quad (27)$$

It appears that the data of $D^+ \rightarrow f_0\pi^+, \sigma\pi^+$ favor the 2-quark picture of the light scalars, while the measurements of $D^0 \rightarrow f_0\pi^0, \sigma\pi^0$ prefer the 4-quark scenario. Moreover, the $D^0 \rightarrow a_0^+K^-$

TABLE VIII: Extracted flavor amplitude parameters from fits to the $D \rightarrow SP$ decays, where S only refers to lighter scalar mesons here. The amplitude magnitudes are in units of 10^{-6} GeV. In these fits, we set $C' = 0$, $E' = E$, and $A' = -A$, with reasons explained in the text. The strong phases $\delta_{E,A}$ are associated with the E and A amplitudes, respectively. For Fits (A) and (B), we take the mixing angle $\theta = 25^\circ$.

Scheme	I		II	
Fit	(A)	(B)	(A)	(B)
$ T $	$2.14^{+0.16}_{-0.15}$	—	1.55 ± 0.07	—
$ A $	$3.16^{+0.10}_{-0.11}$	—	$2.15^{+0.30}_{-0.45}$	—
δ_A	$(31 \pm 2)^\circ$	—	$(35^{+9}_{-11})^\circ$	—
$ C $	—	$1.44^{+0.15}_{-0.14}$	—	$1.90^{+0.36}_{-0.22}$
$ E $	—	$1.20^{+0.05}_{-0.06}$	—	$1.18^{+0.05}_{-0.06}$
δ_E	—	$(168^{+35}_{-10})^\circ$	—	$(152^{+3}_{-2})^\circ$
$\chi^2_{\min} / \text{d.o.f}$	2.61/2	5.07/1	8.29/2	0.74/1

and $a_0^+ \pi^-$ modes will be dominated by the W -exchange diagram, E . The $D^+ \rightarrow a_0^+ \bar{K}^0$ mode is dominated by the C amplitude. The $D^0 \rightarrow \bar{K}_0^{*0} \pi^0$, $D^+ \rightarrow K_0^{*0} \pi^+$, and $D_s^+ \rightarrow K_0^{*0} K^+$ are dominated by the E' , T , and A amplitudes, respectively.

Table VII is divided into two parts separated by double lines. The upper part involves only light scalar mesons (f_0 , a_0 , σ , and κ), whereas the lower part involves the K_0^* mesons in the heavier nonet representation. This division is made because the amplitudes of the same topology in these two groups have no *a priori* relations. We first note that none of the currently measured modes involve the T' amplitude. Secondly, one can simplify the upper part of the table by setting $C' = 0$, for the decay constants of scalar mesons are expected to be either identically zero or relatively small. Moreover, the modes in the lower part of Table VII have the same amplitude decomposition in the two schemes and involve the T , C' , E' , and A amplitudes. One cannot set $C' = 0$ here because the decay constant of K_0^* is non-negligible, as commented after Eq. (14). In this case, there are more theory parameters than observables, barring a fit.

In the following, we will perform two sets of fit [(A) and (B)] to the modes involving only the lighter scalar mesons in the flavor diagram formalism. Fits (A) includes the five measured D^+ and D_s^+ decays. As noted in Eq. (27), the $f_0 \pi^+$ and $\sigma \pi^+$ modes are related because of the same flavor amplitude combination. Here we have to assume a relation between A and A' in order to reduce the number of parameters. Without further theoretical guidance, we have tried the cases $A' = A$ and $A' = -A$ for simplicity and found that the latter renders an equally good or better fit than the former. The strong phase δ_A in Table VIII is measured with respect to T , which is assumed real. We also note here that all the strong phases given in the table are subject to a two-fold ambiguity ($\delta \rightarrow -\delta$). From the $\chi^2_{\min}$ values, one sees that Scheme I fits better than Scheme II in these modes. This is understandable because the 2-quark picture explains better the observed rates of the $f_0 \pi^+$ and $\sigma \pi^+$ decays. In either scheme, $|A|$ is about 1.5 times larger than $|T|$, showing the importance of the W -annihilation contribution. Besides, the extracted relative strong phase is robust.

TABLE IX: Topological amplitudes and branching fractions of $D \rightarrow AP$ decays. The notation is explained in the main text. Theory predictions are made within the factorization approach, with the mixing angle $\theta_{K_1} = 50.8^\circ$.

Decay	Amplitude	Theory	Experiment
$D^+ \rightarrow \bar{K}_1^0(1270)\pi^+$	$V_{cs}^*V_{ud}[(T_A + C'_A)\sin\theta_{K_1} + (T_B + C'_B)\cos\theta_{K_1}]$	4.7×10^{-3}	$< 7 \times 10^{-3}$
$D^+ \rightarrow \bar{K}_1^0(1400)\pi^+$	$V_{cs}^*V_{ud}[(T_A + C'_A)\cos\theta_{K_1} - (T_B + C'_B)\sin\theta_{K_1}]$	2.2%	$(3.8 \pm 1.3)\%$
$D^+ \rightarrow \bar{K}^0 a_1^+(1260)$	$V_{cs}^*V_{ud}(T'_A + C_A)$	8.2%	$(7.0 \pm 1.2)\%$
$D^+ \rightarrow \bar{K}^0 b_1^+(1235)$	$V_{cs}^*V_{ud}(T'_B + C_B)$	2.2×10^{-3}	
$D^0 \rightarrow K_1^-(1270)\pi^+$	$V_{cs}^*V_{ud}[(T_A + E'_A)\sin\theta_{K_1} + (T_B + E'_B)\cos\theta_{K_1}]$	5.2×10^{-3}	$(1.14 \pm 0.32)\%$
$D^0 \rightarrow K_1^-(1400)\pi^+$	$V_{cs}^*V_{ud}[(T_A + E'_A)\cos\theta_{K_1} - (T_B + E'_B)\sin\theta_{K_1}]$	1.4×10^{-4}	$< 1.2\%$
$D^0 \rightarrow \bar{K}_1^0(1270)\pi^0$	$V_{cs}^*V_{ud}\frac{1}{\sqrt{2}}[(C'_A - E'_A)\sin\theta_{K_1} + (C'_B - E'_B)\cos\theta_{K_1}]$	6.6×10^{-3}	
$D^0 \rightarrow \bar{K}_1^0(1400)\pi^0$	$V_{cs}^*V_{ud}\frac{1}{\sqrt{2}}[(C'_A - E'_A)\cos\theta_{K_1} - (C'_B - E'_B)\sin\theta_{K_1}]$	3.2×10^{-3}	$< 3.7\%$
$D^0 \rightarrow K_1^+(1270)K^-$	$V_{cs}^*V_{us}[(T'_A + E_A)\sin\theta_{K_1} + (T'_B + E_B)\cos\theta_{K_1}]$	4.6×10^{-4}	
$D^0 \rightarrow K_1^-(1270)K^+$	$V_{cs}^*V_{us}[(T_A + E'_A)\sin\theta_{K_1} + (T_B + E'_B)\cos\theta_{K_1}]$	8.2×10^{-5}	
$D^0 \rightarrow K_1^\pm(1270)K^\mp$		5.4×10^{-4}	$(8.1 \pm 1.8) \times 10^{-4}$
$D^0 \rightarrow K^- a_1^+(1260)$	$V_{cs}^*V_{ud}(T'_A + E_A)$	2.7%	$(7.9 \pm 1.1)\%$
$D^0 \rightarrow \bar{K}^0 a_1^0(1260)$	$V_{cs}^*V_{ud}\frac{1}{\sqrt{2}}(C_A - E_A)$	1.2×10^{-4}	$< 1.9\%$
$D^0 \rightarrow \pi^- a_1^+(1260)$	$V_{cd}^*V_{ud}(T'_A + E_A)$	5.1×10^{-3}	$(8.98 \pm 0.62) \times 10^{-3}$
$D^0 \rightarrow K^- b_1^+(1235)$	$V_{cs}^*V_{ud}(T'_B + E_B)$	1.7×10^{-5}	
$D^0 \rightarrow \bar{K}^0 b_1^0(1235)$	$V_{cs}^*V_{ud}\frac{1}{\sqrt{2}}(C_B - E_B)$	3.0×10^{-4}	

Fit (B) includes the four measured D^0 decays. Here it does not matter what relation we assume between E and E' , as far as the $\chi^2_{\min}$ value is concerned. The only effect is on the size and phase of the E amplitude. This is because the $f_0\pi^0$ and $\sigma\pi^0$ modes are related by the same flavor amplitude combination, as also noted in Eq. (27). We assume $E' = E$ for an explicit fit. In Table VIII, the strong phase δ_E is measured with respect to C , which is assumed real. The $\chi^2_{\min}$ values show that Scheme II explains this set of data better, as noted below Eq. (27). It is worth noting that the magnitudes and relative phase extracted in either scheme are roughly the same. Also, the amplitudes C and E are almost opposite in phase, as required primarily by the $D^0 \rightarrow a_0^0 \bar{K}^0$ decay. Finally, based on the $\chi^2_{\min}$ values of these fits, the current data still cannot differentiate the two schemes yet.

B. $D \rightarrow AP$

The topological amplitudes for $D \rightarrow AP$ decays are given in Table IX. Instead of using subscripts of A and P to complicate the notation, we use the primed (unprimed) amplitudes to indicate that the spectator quark in the D meson ends up in the pseudoscalar (axial-vector) meson in the final state. The subscripts A and B refer to the amplitudes associated with the 3P_1 and 1P_1 axial-vector mesons, respectively. However, as we will see later in Section VB, the factorization approach predicts that such a distinction is only necessary for the T amplitudes. The assump-

TABLE X: Topological amplitudes and branching fractions of $D \rightarrow TP$ decays. The notation is explained in the main text. Theory predictions are made within the factorization approach, with the mixing angle $\theta_{f_2} = 7.8^\circ$.

Decay	Amplitude	Theory	Experiment
$D^+ \rightarrow f_2 \pi^+$	$\frac{1}{\sqrt{2}} V_{cd}^* V_{ud} \cos \theta_{f_2} (T + C' + A + A') + V_{cs}^* V_{us} \sin \theta_{f_2} C'$	0.9×10^{-6}	$(8.8 \pm 1.6) \times 10^{-4}$
$D^+ \rightarrow a_2^+ \bar{K}^0$	$V_{cs}^* V_{ud} (T' + C)$	4.5×10^{-7}	$< 3.0 \times 10^{-3}$
$D^+ \rightarrow \bar{K}_2^{*0} \pi^+$	$V_{cs}^* V_{ud} (T + C')$	1.9×10^{-5}	$(6.3 \pm 1.2) \times 10^{-4}$
$D^+ \rightarrow K_2^{*0} \pi^+$	$V_{cd}^* V_{us} (C' + A)$	0	$(1.5 \pm 1.0) \times 10^{-4}$
$D^0 \rightarrow f_2 \pi^0$	$\frac{1}{2} V_{cd}^* V_{ud} \cos \theta_{f_2} (C' - C - E' - E) + \frac{1}{\sqrt{2}} V_{cs}^* V_{us} \sin \theta_{f_2} C'$	5.1×10^{-8}	$(3.4 \pm 0.4) \times 10^{-4}$
$D^0 \rightarrow f_2 \bar{K}^0$	$V_{cs}^* V_{ud} \left[\frac{1}{\sqrt{2}} \cos \theta_{f_2} (C + E) + \sin \theta_{f_2} E' \right]$	1.5×10^{-7}	$(5.0^{+3.5}_{-2.1}) \times 10^{-4}$
$D^0 \rightarrow a_2^+ K^-$	$V_{cs}^* V_{ud} (T' + E)$	0	$< 2 \times 10^{-3}$
$D^0 \rightarrow K_2^{*-} \pi^+$	$V_{cs}^* V_{ud} (T + E')$	7.5×10^{-6}	$(2.1^{+1.2}_{-0.7}) \times 10^{-3}$
$D_s^+ \rightarrow f_2 \pi^+$	$V_{cs}^* V_{ud} \left[\frac{1}{\sqrt{2}} \cos \theta_{f_2} (A + A') + \sin \theta_{f_2} T \right]$	7.0×10^{-6}	$(1.9 \pm 0.4) \times 10^{-3}$
$D_s^+ \rightarrow K_2^{*0} \pi^+$	$V_{cd}^* V_{ud} T + V_{cs}^* V_{us} A$	2.4×10^{-6}	$(1.5 \pm 1.2) \times 10^{-3}$

tion of $C_A = C_B$ and $E_A = E_B$ can be checked by comparing the rates of $D^0 \rightarrow \bar{K}^0 a_1^0(1260)$ and $D^0 \rightarrow \bar{K}^0 b_1^0(1235)$, which are seen to be roughly the same up to a tiny phase space difference. With the flavor symmetry assumption, the magnitudes of the invariant amplitudes of $D^0 \rightarrow \pi^- a_1^+(1260)$ and $D^0 \rightarrow K^- a_1^+(1260)$ should differ by a factor of $\lambda \simeq 0.2253$, which is to be compared with 0.171 ± 0.013 given by the current data. A distinctive feature between the Cabibbo-allowed D^0 and D^+ decays is that the W -exchange diagrams (E) only involve in the former.

Current data for $D \rightarrow AP$ decays (only six branching fractions) are still insufficient for a sensible fit. The theory predictions in Table IX are based on the factorization calculations given in Section VB.

C. $D \rightarrow TP$

The topological amplitudes for $D \rightarrow TP$ decays are given in Table X. Here we also use the unprimed (primed) symbols to indicate that the spectator quark of the D meson ends up in the even-parity (tensor) and odd-parity (pseudoscalar) mesons in the final state, respectively. There should be no confusion even though the amplitude symbols used here are identical to those in the $D \rightarrow SP$ case.

As described before, the decay constants of the tensor mesons vanish identically. Therefore, one can set $T' = C' = 0$ in Table X. The decay $D^+ \rightarrow \bar{K}_2^{*0} K^+$ is kinematically forbidden as the K_2^* mass is above the kinematic threshold, though it is physically allowed through the width of K_2^* . We therefore will not include it in our fit. Note that the measured D^+ and D_s^+ decays only involve the T , A , and A' amplitudes. The results of fits to these decay modes are given in Table XI. Since

TABLE XI: Extracted flavor amplitude parameters from fits to the $D \rightarrow TP$ decays. The amplitude magnitudes are in units of 10^{-6}GeV^{-1} , the same as M in Eq. (26). In these fits, we set $C' = 0$ and $\theta_{f_2} = 7.8^\circ$. The strong phases $\delta_{A,A'}$ are associated with the A and A' amplitudes, respectively, relative to T . The contents of different fits are described in the main text.

Parameter	$ T $	$ A $	δ_A	$ A' $	$\delta_{A'}$	$\chi^2_{\min} / \text{d.o.f.}$
Fit (A)	$1.85^{+0.14}_{-0.16}$	$12.63^{+0.15}_{-0.16}$	$(0 \pm 3)^\circ$	$11.24^{+0.16}_{-0.15}$	$(180 \pm 3)^\circ$	11.4/0
Fit (B)	$8.13^{+0.39}_{-0.42}$	1.59 ± 0.08	$(173^{+26}_{-12})^\circ$	—	—	0/0

there are only three measured modes in the D^0 decays whereas at least four additional parameters have to be introduced even if we set $E' = E$, it is impossible to determine the magnitudes and strong phases of C and E by considering a global fit to the TP decays. For that, a determination of $\mathcal{B}(D^0 \rightarrow a_2^+ K^-)$ is crucial.

Fit (A) in Table XI includes all the available D^+ and D_s^+ decay modes except for $D^+ \rightarrow \overline{K}_2^{*0} K^+$; there are thus 5 observables for 5 parameters. According to the results of Fit (A), the current data favor relatively large W -annihilation diagrams. This is because $|A|$ and $|A'|$ are constrained respectively by the doubly Cabibbo-suppressed $D^+ \rightarrow K_2^{*0} \pi^+$ mode and the singly Cabibbo-suppressed $D_s^+ \rightarrow f_2 K^+$ mode to be large. Moreover, A and A' are about the same size but opposite in phase so that the branching fraction of $D_s^+ \rightarrow f_2 \pi^+$ falls in the ball park. On the other hand, $|T|$ is largely constrained by the Cabibbo-favored $D^+ \rightarrow \overline{K}_2^{*0} \pi^+$ mode to be small. The largest contribution in the $\chi^2_{\min}$ value comes from the $D^+ \rightarrow f_2 \pi^+$ mode (~ 10.1). This is a manifestation of the disparity between the Cabibbo-favored $D_s^+ \rightarrow f_2 \pi^+$ decay and the singly Cabibbo-suppressed $D^+ \rightarrow f_2 \pi^+$ decay that are seen to have similar branching ratios, if $|T|$ is constrained not to play a role here.

In view of the possibly problematic $D^+ \rightarrow K_2^{*0} \pi^+$ and $\overline{K}_2^{*0} \pi^+$ modes, we exclude them in Fit (B) and set $A' = A$ for simplicity; there are then 3 observables for 3 parameters. We note that it is not illuminating to consider $A' = -A$ here because two of these modes involve the combination $A + A'$ as the major contribution. In Fit (B), $|T|$ becomes larger and A much suppressed. Also, the relative strong phase between A and T is almost opposite to that in Fit (A).

The third column in Table X lists theory predictions based on factorization assumption to be discussed in Section V C. A comparison between the predictions and the measured values shows an apparent deficit in theory account of the decay amplitudes. First, the magnitude of the tree contribution in the factorization approach is even smaller than the value of $|T|$ in Table XI. Secondly, it is necessary to invoke the annihilation type of amplitudes to explain the observed data.

V. FACTORIZATION APPROACH

The diagrammatic approach has been applied quite successfully to hadronic decays of charmed mesons into PP and VP final states [2, 6, 67–72]. When generalized to the decay modes involving an even-parity light meson in the final state, it appears that the current data are still insufficient

for us to fully extract the information of all amplitudes. Moreover, as shown in Tables VIII and XI, the extracted parameters do not present a coherent picture yet. Therefore, we take the naive factorization formalism as a complementary approach to estimate the rates of these decay modes. In this framework, the W -exchange and -annihilation type of cotributions will be neglected. We discuss the three categories of decays in the following subsections separately.

A. $D \rightarrow SP$

The factorizable amplitudes for the $D \rightarrow SP$ decays involve

$$\begin{aligned} X^{(DS,P)} &= \langle P(q)|(V-A)_\mu|0\rangle\langle S(p)|(V-A)^\mu|D(p_D)\rangle, \\ X^{(DP,S)} &= \langle S(q)|(V-A)_\mu|0\rangle\langle P(p)|(V-A)^\mu|D(p_D)\rangle, \end{aligned} \quad (28)$$

with the expressions

$$X^{(DS,P)} = -f_P(m_D^2 - m_S^2)F_0^{DS}(q^2), \quad X^{(DP,S)} = f_S(m_D^2 - m_P^2)F_1^{DP}(q^2), \quad (29)$$

where use of Eqs. (12) and (15) has been made. The decay amplitudes of $D \rightarrow K_0^*P$ thus read

$$\begin{aligned} A(D^+ \rightarrow \bar{K}_0^{*0}\pi^+) &= \frac{G_F}{\sqrt{2}}V_{cs}^*V_{ud}\left[-a_1f_\pi(m_D^2 - m_{K_0^*}^2)F_0^{DK_0^*}(m_\pi^2) \right. \\ &\quad \left. + a_2f_{K_0^*}(m_D^2 - m_\pi^2)F_0^{D\pi}(m_{K_0^*}^2)\right], \\ A(D^0 \rightarrow K_0^{*-}\pi^+) &= -\frac{G_F}{\sqrt{2}}V_{cs}^*V_{ud}a_1f_\pi(m_D^2 - m_{K_0^*}^2)F_0^{DK_0^*}(m_\pi^2), \\ A(D^0 \rightarrow \bar{K}_0^{*0}\pi^0) &= \frac{G_F}{2}V_{cs}^*V_{ud}a_2f_{K_0^*}(m_D^2 - m_\pi^2)F_0^{D\pi}(m_{K_0^*}^2), \\ A(D_s^+ \rightarrow K_0^{*0}\pi^+) &= -\frac{G_F}{\sqrt{2}}a_1V_{cd}^*V_{ud}f_\pi(m_{D_s}^2 - m_{K_0^*}^2)F_0^{D_sK_0^*}(m_\pi^2), \end{aligned} \quad (30)$$

and likewise for the other $D \rightarrow SP$ decays.

Using the decay constants and form factors given in Section III A, the predicted rates of $(D, D_s) \rightarrow (f_0, K_0^*)P$ are computed and listed in Table XII, where we have used $a_1 = 1.22$, $a_2 = -0.66$ and taken the form factors for D to π and K transitions from the recent CLEO-c measurements of semileptonic D meson decays to π and K mesons [73]. In order to test the factorization approach, we should focus on the modes in which weak annihilations (W -exchange or W -annihilation) are absent or suppressed. The Cabibbo-allowed decays $D^+ \rightarrow \bar{K}_0^{*0}\pi^+$ and $D_s^+ \rightarrow f_0\pi^+$ satisfy this criterion: the weak annihilation amplitude is absent in the former and suppressed by the $f_0 - \sigma$ mixing in the latter. We see from Table XII that factorization works well for these two modes. For Cabibbo-allowed $D^+ \rightarrow PP$ or VP decays, it is known that the color-allowed T and color-suppressed C amplitudes interfere destructively due to the opposite sign of the parameters a_1 and a_2 . However, it is the other way around for Cabibbo-allowed $D^+ \rightarrow SP$ and AP decays. From Eq. (30), it is obvious that the a_1 and a_2 terms in the decay amplitude of $D^+ \rightarrow \bar{K}_0^{*0}\pi^+$ interfere constructively. If they interfered destructively, one would have $\mathcal{B}(D^+ \rightarrow \bar{K}_0^{*0}\pi^+) = 5.4 \times 10^{-5}$ which is too small compared to experiment. Numerically, we obtain $|T| = 9.2 \times 10^{-7}$ GeV and $|C'| = 1.0 \times 10^{-6}$ GeV for this mode. Therefore, even though C' is

TABLE XII: The predicted branching fractions for various $D \rightarrow SP$ decays with the scalar mesons treated as $q\bar{q}$ ground states. For simplicity, we have dropped the mass identification for $f_0(980)$ and $K_0^*(1430)$. The $f_0 - \sigma$ mixing angle θ is taken to be 25° . Theory predictions are made within the factorization approach in which the weak annihilation topologies (E and A) are neglected.

Decay	Amplitude	$\mathcal{B}_{\text{theory}}$	$\mathcal{B}_{\text{expt}}$
$D^0 \rightarrow f_0 \pi^0$	$\frac{1}{2} V_{cd}^* V_{ud} (-C + C' - E - E') \sin \theta$ $+ \frac{1}{\sqrt{2}} V_{cs}^* V_{us} C' \cos \theta$	7.8×10^{-6}	$(1.0 \pm 0.3) \times 10^{-4}$
$\rightarrow f_0 \bar{K}^0$	$V_{cs}^* V_{ud} [\frac{1}{\sqrt{2}} (C + E) \sin \theta + E' \cos \theta]$	3.5×10^{-4}	$(8.0_{-2.2}^{+2.5}) \times 10^{-3}$
$D^+ \rightarrow f_0 \pi^+$	$\frac{1}{\sqrt{2}} V_{cd}^* V_{ud} (T + C' + A + A') \sin \theta$ $+ V_{cs}^* V_{us} C' \cos \theta$	1.4×10^{-4}	$(4.5 \pm 1.4) \times 10^{-4}$
$\rightarrow f_0 K^+$	$V_{cd}^* V_{us} [\frac{1}{\sqrt{2}} (T + A') \sin \theta + A \cos \theta]$	1.1×10^{-5}	$(1.6 \pm 1.0) \times 10^{-4}$
$D_s^+ \rightarrow f_0 \pi^+$	$V_{cs}^* V_{ud} (T \cos \theta + (A + A') \sin \theta / \sqrt{2})$	1.3%	$(1.8 \pm 0.5)\%$
$D^0 \rightarrow K_0^{*-} \pi^+$	$V_{cs}^* V_{ud} (T + E')$	2.0×10^{-3}	$(8.2 \pm 1.4) \times 10^{-3}$
$\rightarrow \bar{K}_0^{*0} \pi^0$	$V_{cs}^* V_{ud} (C' - E') / \sqrt{2}$	1.2×10^{-3}	$(9.2_{-2.6}^{+8.1}) \times 10^{-3}$
$D^+ \rightarrow \bar{K}_0^{*0} \pi^+$	$V_{cs}^* V_{ud} (T + C')$	2.3%	$(1.8 \pm 0.3)\%$ ^a
$D_s^+ \rightarrow \bar{K}_0^{*0} K^+$	$V_{cs}^* V_{ud} (C' + A)$	5.5×10^{-4}	$(3.6 \pm 0.7) \times 10^{-3}$
$\rightarrow K_0^{*0} \pi^+$	$V_{cd}^* V_{ud} T + V_{cs} V_{us}^* A$	2.6×10^{-4}	$(8.1 \pm 6.5) \times 10^{-4}$

^aData from E791 [31]; see also Eq. (5).

suppressed by the smallness of $f_{K_0^*(1430)}$ and a_2 , it is enhanced sizably by the mass squared term ($m_D^2 - m_\pi^2$) and the form factor $F_0^{D\pi}$ at $q^2 = m_{K_0^*}^2$.

From Table XII we see that the predicted rates for the other $D \rightarrow \bar{K}_0^* P$ ($P = \pi, K$) decays are smaller than experiments by a factor of $2 \sim 8$. Note that they always receive weak annihilation contributions (E or A). Under the factorization hypothesis, the factorizable W -exchange and W -annihilation amplitudes are suppressed due to the smallness of the form factor at large $q^2 = m_D^2$. This corresponds to the so-called helicity suppression. However, sizable long-distance weak annihilation can be induced via FSIs. For charm decays, it is expected that the long-distance weak annihilation is dominated by resonant FSIs. That is, the FSI via $q\bar{q}$ resonances is usually the most important one due to the fact that an abundant spectrum of resonances is known to exist at energies close to the masses of the charmed mesons. The diagrammatic-approach analysis in the last section suggests that weak annihilation diagrams are comparable to or even larger than the color-allowed tree amplitude T . Therefore, it is conceivable that the inclusion of E' and A amplitudes can account for $\mathcal{B}(D \rightarrow \bar{K}_0^* P)$. For $D(D_s^+) \rightarrow f_0 \pi$ and $f_0 K$ decays, the calculated branching fractions are typically too small by about one order of magnitude. To enhance the rates in this case, the weak annihilation contributions have to be larger than the color-allowed tree amplitude, as shown in Table VIII. Such an amplitude hierarchy poises a difficulty in theoretical understanding. While the non-strange content of $f_0(980)$ is small in the two-quark model for light scalars, it is not so in the tetraquark picture. This suggests that one should treat the light scalar mesons as bound states of $qq\bar{q}\bar{q}$. Unfortunately, the naive quark model is not applicable to evaluating the form factors for

the transition of D to a 4-quark state.

B. $D \rightarrow AP$

The factorizable amplitudes for the $D \rightarrow AP$ decays involve

$$\begin{aligned} X^{(DA,P)} &= \langle P(q)|(V-A)_\mu|0\rangle\langle A(p)|(V-A)^\mu|D(p_D)\rangle, \\ X^{(DP,A)} &= \langle A(q)|(V-A)_\mu|0\rangle\langle P(p)|(V-A)^\mu|D(p_D)\rangle. \end{aligned} \quad (31)$$

with the expressions

$$X^{(DA,P)} = 2f_P m_A V_0^{DA}(q^2)(\epsilon^* \cdot p_D), \quad X^{(DP,A)} = -2f_A m_A F_1^{DP}(q^2)(\epsilon^* \cdot p_D). \quad (32)$$

It is then straightforward to write down the factorizable amplitudes of $D \rightarrow K_1(1270)\pi$ and $D \rightarrow K_1(1400)\pi$ decays (dropping the overall $\epsilon^* \cdot p_D$ terms for simplicity):

$$\begin{aligned} A(D^+ \rightarrow \bar{K}_1^0(1270)\pi^+) &= \frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} \left[2a_1 m_{K_1(1270)} f_\pi (\sin \theta_{K_1} V_0^{DK_{1A}}(m_\pi^2) + \cos \theta_{K_1} V_0^{DK_{1B}}(m_\pi^2)) \right. \\ &\quad \left. - 2a_2 m_{K_1(1270)} f_{K_1(1270)} F_1^{D\pi}(m_{K_1(1270)}^2) \right], \\ A(D^+ \rightarrow \bar{K}_1^0(1400)\pi^+) &= \frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} \left[2a_1 m_{K_1(1400)} f_\pi (\cos \theta_{K_1} V_0^{DK_{1A}}(m_\pi^2) - \sin \theta_{K_1} V_0^{DK_{1B}}(m_\pi^2)) \right. \\ &\quad \left. - 2a_2 m_{K_1(1400)} f_{K_1(1400)} F_1^{D\pi}(m_{K_1(1400)}^2) \right], \\ A(D^0 \rightarrow K_1^-(1270)\pi^+) &= \frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} \left[2a_1 m_{K_1(1270)} f_\pi (\sin \theta_{K_1} V_0^{DK_{1A}}(m_\pi^2) + \cos \theta_{K_1} V_0^{DK_{1B}}(m_\pi^2)) \right], \\ A(D^0 \rightarrow K_1^-(1400)\pi^+) &= \frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} \left[2a_1 m_{K_1(1400)} f_\pi (\cos \theta_{K_1} V_0^{DK_{1A}}(m_\pi^2) - \sin \theta_{K_1} V_0^{DK_{1B}}(m_\pi^2)) \right], \\ A(D^0 \rightarrow \bar{K}_1^0(1270)\pi^0) &= -\frac{G_F}{2} V_{cs}^* V_{ud} \left[2a_2 m_{K_1(1270)} f_{K_1(1270)} F_1^{D\pi}(m_{K_1(1270)}^2) \right], \\ A(D^0 \rightarrow \bar{K}_1^0(1400)\pi^0) &= -\frac{G_F}{2} V_{cs}^* V_{ud} \left[2a_2 m_{K_1(1400)} f_{K_1(1400)} F_1^{D\pi}(m_{K_1(1400)}^2) \right], \\ A(D^0 \rightarrow K_1^+(1270)K^-) &= -\frac{G_F}{\sqrt{2}} V_{cs}^* V_{us} \left[2a_1 m_{K_1(1270)} f_{K_1(1270)} F_1^{DK}(m_{K_1(1270)}^2) \right], \\ A(D^0 \rightarrow K_1^-(1270)K^+) &= \frac{G_F}{\sqrt{2}} V_{cs}^* V_{us} \left[2a_1 m_{K_1(1270)} f_K (\sin \theta_{K_1} V_0^{DK_{1A}}(m_K^2) + \cos \theta_{K_1} V_0^{DK_{1B}}(m_K^2)) \right], \end{aligned} \quad (33)$$

where we have taken into account the $K_{1A} - K_{1B}$ mixing given by Eq. (17) and neglected the short-distance factorizable W -exchange contributions. Likewise, the $D \rightarrow Ka_1(1260)$ and $D \rightarrow Kb_1(1235)$ decay amplitudes read

$$\begin{aligned} A(D^+ \rightarrow \bar{K}^0 a_1^+(1260)) &= -\frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} \left[2a_1 f_{a_1} m_{a_1} F_1^{DK}(m_{a_1}^2) - 2a_2 f_K m_{a_1} V_0^{Da_1}(m_K^2) \right], \\ A(D^0 \rightarrow K^- a_1^+(1260)) &= -\frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} 2a_1 f_{a_1} m_{a_1} F_1^{DK}(m_{a_1}^2), \\ A(D^0 \rightarrow \bar{K}^0 a_1^0(1260)) &= \frac{G_F}{2} V_{cs}^* V_{ud} 2a_2 f_K m_{a_1} V_0^{Da_1}(m_K^2), \\ A(D^0 \rightarrow \pi^- a_1^+(1260)) &= -\frac{G_F}{\sqrt{2}} V_{cd}^* V_{ud} 2a_1 f_{a_1} m_{a_1} F_1^{D\pi}(m_{a_1}^2), \\ A(D^+ \rightarrow \bar{K}^0 b_1^+(1235)) &= -\frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} \left[2a_1 f_{b_1} m_{b_1} F_1^{DK}(m_{b_1}^2) - 2a_2 f_K m_{b_1} V_0^{Db_1}(m_K^2) \right], \end{aligned}$$

$$\begin{aligned}
A(D^0 \rightarrow K^- b_1^+(1235)) &= -\frac{G_F}{\sqrt{2}} V_{cs}^* V_{ud} 2a_1 f_{b_1} m_{b_1} F_1^{DK}(m_{b_1}^2) , \\
A(D^0 \rightarrow \bar{K}^0 b_1^0(1235)) &= \frac{G_F}{2} V_{cs}^* V_{ud} 2a_2 f_K m_{b_1} V_0^{D b_1}(m_K^2) .
\end{aligned} \tag{34}$$

Using the decay constants and form factors presented in Section III B, the predicted rates of $D \rightarrow \bar{K}_1 \pi, \bar{K} a_1, \bar{K} b_1$ decays are listed in Table IX. To test the validity of the factorization hypothesis, we focus on the D^+ decay to $\bar{K}_1 \pi^+$ and $\bar{K}^0 a_1(1260)$ which are free of contamination from weak annihilations. We see that the predictions are in agreement with experiment for these D^+ decays and hence factorization works for $D \rightarrow AP$, just as the case of $D \rightarrow SP$. The predicted rates for D^0 decays are slightly smaller, which implies the importance of the W -exchange contribution to D^0 decay modes. The theoretical calculations presented in Table IX are for the $K_1(1270)$ - $K_1(1400)$ mixing angle $\theta_{K_1} = 50.8^\circ$. When the other solution $\theta_{K_1} = -44.8^\circ$ is used, we find the predictions $\mathcal{B}(D^+ \rightarrow \bar{K}_1^0(1270)\pi^+) = 2.9\%$, $\mathcal{B}(D^+ \rightarrow \bar{K}_1^0(1400)\pi^+) = 2.0 \times 10^{-3}$ and $\mathcal{B}(D^0 \rightarrow K_1^+(1270)\pi^+) = 9.1 \times 10^{-5}$, all in sharp disagreement with the data. Historically, it was first pointed out in Ref. [55] that a negative mixing angle θ_{K_1} is ruled out by the data of $D^+ \rightarrow \bar{K}_1^0(1270)\pi^+$ and $D^0 \rightarrow K_1^-(1270)\pi^+$.⁶ It was realized later that the negative θ_{K_1} solution is also ruled out by the experimental measurements of $B \rightarrow K_1(1270)\gamma$ and $B \rightarrow K_1(1400)\gamma$ [74].

It is of interest to notice that the $D^0 \rightarrow K_1^\pm(1400)K^\mp$ decay is not kinematically allowed, yet a branching fraction comparable to $\mathcal{B}(D^0 \rightarrow K_1^\pm(1270)K^\mp \rightarrow K^\mp K^\pm \pi^- \pi^+)$ has been observed. Since the width of $K_1(1400)$ is 174 ± 13 MeV, the $D^0 \rightarrow K_1^\pm(1400)K^\mp$ decay followed by $K_1^\pm(1400) \rightarrow K^\pm \pi^+ \pi^-$ is certainly allowed.

C. $D \rightarrow TP$

Since the decay constant of tensor meson vanishes, the factorizable amplitude of $D \rightarrow TP$ always involves the expression

$$\begin{aligned}
X^{(DT,P)} &= \langle P(q)|(V-A)_\mu|0\rangle \langle T(p)|(V-A)^\mu|D(p_D)\rangle \\
&= i f_P \varepsilon_{\mu\nu}^* p_D^\mu p_D^\nu \left[k(m_P^2) + b_+(m_P^2)(m_D^2 - m_T^2) + b_-(m_P^2)m_P^2 \right] ,
\end{aligned} \tag{35}$$

where use has been made of Eq. (24). The decay rate is given by Eq. (26). In general, TP final states are suppressed relative to PP states due to less available phase space. More precisely,

$$\frac{\Gamma(D \rightarrow TP)}{\Gamma(D \rightarrow P_1 P_2)} = \frac{2}{3} \frac{p_T^5}{p_P} \left(\frac{m_D}{m_T} \right)^4 \left| \frac{M(D \rightarrow TP)}{M(D \rightarrow P_1 P_2)} \right|^2 , \tag{36}$$

where we have changed the notation slightly so that p_T denotes the c. m. momentum of the tensor meson and p_P is the c. m. momentum of the pseudoscalar meson P_1 or P_2 in the charmed meson rest frame. The kinematic factor $\frac{2}{3}(p_T^5/p_P)(m_D/m_T)^4$ is typically of order $(1-4) \times 10^{-2} \text{ GeV}^{-4}$. An inspection of Table X indicates that, in the absence of weak annihilation contributions, the Cabibbo-allowed decays $D^+ \rightarrow \bar{K}_2^{*0} \pi^+$ and $D^0 \rightarrow K_2^{*-} \pi^+$ should have the largest decay rates as

⁶ The calculation of [55] was performed in the ISGW2 model [52] which has the opposite sign convention to the CLF model.

they proceed through the color-allowed tree diagram T . It is easily seen that all other W -emission amplitudes in $D \rightarrow a_2 \bar{K}$, $D \rightarrow f_2 \pi$ and $D \rightarrow f_2 \bar{K}$ are suppressed for various reasons. For example, it is suppressed by the vanishing decay constant of the tensor meson, by the small $f_2 - f'_2$ mixing angle, by the parameter a_2 , or by the Cabibbo angle.

From Table X we see that the predicted branching ratio of $D^+ \rightarrow \bar{K}_2^{*0} \pi^+$ is of order 10^{-5} , which is about two orders of magnitude smaller than experiment. Indeed, the theoretical calculation gives $|T| = 3 \times 10^{-7}$ in the unit of GeV^{-1} , which is much smaller than the value of $|T|$ listed in Table XI. As for the decay $D^0 \rightarrow K_2^{*-} \pi^+$, its rate is similar to that of $D^+ \rightarrow \bar{K}_2^{*0} \pi^+$ but receives an additional W -exchange contribution. A fit of this mode to experiment will require $|E| > |T|$, namely, W -exchange dominates over the external W -emission. The current measurement of $\mathcal{B}(D^+ \rightarrow \bar{K}_2^{*0} K^+)$ is problematic as it is Cabibbo-suppressed and yet the measured rate is larger than $\mathcal{B}(D^+ \rightarrow \bar{K}_2^{*0} \pi^+)$.

All the predictions shown in Table X are too small by at least two orders of magnitude, as originally noticed in Ref. [27]. In order to resolve the enormous discrepancy between theory and experiment for TP modes, one may consider possible form factor enhancement and finite width effects. One may compare the $D \rightarrow T$ transition form factors calculated in different models: the relativistic light-front quark model (see Table V) and the ISGW and ISGW2 quark models (see Table II of Ref. [27]). Since the form factors obtained from different models are of the same order, it is very unlikely that they can be enhanced by one order of magnitude to ameliorate the discrepancy. The finite width effect of the tensor resonances will be discussed in the next section.

VI. FINITE WIDTH EFFECTS

Normally we apply the narrow width approximation to extract the two-body branching fraction $\mathcal{B}(D \rightarrow MP)$ from 3-body decay data with M standing for an even-parity meson. There are three cases where the narrow width approximation is not valid or justified and the finite width of the resonance has to be taken into account: (i) The decay $D \rightarrow MP$ is not kinematically allowed. For example, $D^+ \rightarrow \bar{K}_0^{*0} K^+$, $\bar{K}_2^{*0} K^+$ and $D^0 \rightarrow a_0^+(1450) K^-$, $K_1^+(1400) K^-$ are forbidden if the scalar resonances are very narrow and on their mass shells. (ii) The resonance width is not negligible. For example, the widths of σ and κ are very broad, of order $600 - 1000$ and 550 ± 34 MeV, respectively [29]. (iii) The strong decay of resonance is marginally allowed or even forbidden kinematically. For instance, the central values of the $f_0(980)$ and $a_0(980)$ masses are below the threshold for decaying into a charged kaon pair.

In general, the rate of the three-body decay $D \rightarrow P_1 P_2 P$ is given by Ref. [75]⁷

$$\begin{aligned} \Gamma(D \rightarrow SP \rightarrow P_1 P_2 P) &= \frac{1}{2m_D} \int_{(m_1+m_2)^2}^{(m_D-m_P)^2} \frac{dq^2}{2\pi} |M(D \rightarrow SP)|^2 \frac{\lambda^{1/2}(m_D^2, q^2, m_P^2)}{8\pi m_D^2} \\ &\times \frac{1}{(q^2 - m_S^2)^2 + (\Gamma_{12}(q^2)m_S)^2} g_{SP_1 P_2}^2 \frac{\lambda^{1/2}(q^2, m_1^2, m_2^2)}{8\pi q^2} \end{aligned} \quad (37)$$

⁷ The case for $D \rightarrow AP \rightarrow VP_1 P \rightarrow P_1 P_2 P_3 P$ is more complicated and has been discussed in Ref. [22]. The formula for $\Gamma(D \rightarrow TP \rightarrow P_1 P_2 P)$ given in Ref. [27] was erroneous and it is corrected here.

via a scalar resonance, and

$$\begin{aligned} \Gamma(D \rightarrow TP \rightarrow P_1 P_2 P) &= \frac{1}{\pi} \int_{(m_1+m_2)^2}^{(m_D-m_P)^2} \frac{dq^2}{2\pi} |M(D \rightarrow TP)|^2 \frac{p(q^2)^5}{12\pi} \frac{m_D^2}{q^2} \\ &\times \frac{1}{(q^2 - m_T^2)^2 + (\Gamma_{12}(q^2)m_T)^2} g_{TP_1P_2}^2 \frac{p'(q^2)^5}{15\pi q^5} \end{aligned} \quad (38)$$

via a tensor resonance, where λ is the usual triangular function $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2bc - 2ca$, m_1 (m_2) is the mass of P_1 (P_2), $p(q^2) = \lambda^{1/2}(m_D^2, q^2, m_P^2)/(2m_D)$, $p'(q^2) = \lambda^{1/2}(q^2, m_1^2, m_2^2)/(2\sqrt{q^2})$, and $g_{MP_1P_2}$ is the strong coupling to be defined below. The “running” or “comoving” width $\Gamma_{12}(q^2)$ is a function of the invariant mass $m_{12} = \sqrt{q^2}$ of the $P_1 P_2$ system and has the expression [76]

$$\Gamma_{12}(q^2) = \begin{cases} \Gamma_T \frac{m_T}{m_{12}} \left(\frac{p'(q^2)}{p'(m_T^2)} \right)^5 \frac{9+3R^2 p'^2(m_T^2)+R^4 p'^4(m_T^2)}{9+3R^2 p'^2(q^2)+R^4 p'^4(q^2)}, & \text{for } M = T, \\ \Gamma_S \frac{m_S}{m_{12}} \frac{p'(q^2)}{p'(m_S^2)} & \text{for } M = S. \end{cases} \quad (39)$$

Note that the propagator of the resonance has been assumed to be of the Breit-Wigner form. From the measured widths of the resonances, one can determine their strong couplings

$$\Gamma(S \rightarrow P_1 P_2) = g_{SP_1P_2}^2 \frac{p_c}{8\pi m_S^2}, \quad \Gamma(T \rightarrow P_1 P_2) = \frac{g_{TP_1P_2}^2 m_T}{15\pi} \left(\frac{p_c}{m_T} \right)^5. \quad (40)$$

When the resonance width Γ_M is narrow, the expression of the resonant decay rate can be simplified by applying the so-called narrow width approximation

$$\frac{1}{(q^2 - m_M^2)^2 + m_M^2 \Gamma_M^2(q^2)} \approx \frac{\pi}{m_M \Gamma_M} \delta(q^2 - m_M^2). \quad (41)$$

It is easily seen that this leads to the factorization relation Eq. (1) for the resonant three-body decay.

In the following, we illustrate the finite width effects with a few examples.

$$\underline{D^+ \rightarrow \bar{K}_0^{*0} K^+ \rightarrow K^+ K^- \pi^+}$$

With a width of 270 ± 80 MeV for $K_0^*(1430)$, the decay $D^+ \rightarrow \bar{K}_0^{*0} K^+$ followed by $\bar{K}_0^{*0} \rightarrow K^+ K^- \pi^+$ is now physically allowed. In this case one should evaluate the two-step process $\Gamma(D^+ \rightarrow \bar{K}_0^{*0} K^+ \rightarrow K^+ K^- \pi^+)$ and compare the resonant three-body rate with experiment. Using Eq. (37) and assuming that the coupling $g_{SP_1P_2}$ is insensitive to the variation in q^2 when the resonance is off its mass shell, we obtain

$$\mathcal{B}(D^+ \rightarrow \bar{K}_0^{*0} K^+ \rightarrow K^+ K^- \pi^+) = (1.3_{-0.3}^{+0.1}) \times 10^{-4}. \quad (42)$$

This is one order of magnitude smaller than the experimental value, $(1.83 \pm 0.35) \times 10^{-3}$ (see Table I). Since

$$A(D^+ \rightarrow \bar{K}_0^{*0} K^+) = V_{cs}^* V_{us} T + V_{cd}^* V_{ud} A, \quad (43)$$

the inclusion of W -annihilation A will improve the discrepancy between theory and experiment.

$$\underline{D^+ \rightarrow \sigma \pi^+}$$

Since the width of the σ resonance $\Gamma_\sigma = 600 - 1000$ MeV is of the same order of magnitude as its mass, it is important to see its effect on the extraction of the branching fraction $\mathcal{B}(D^+ \rightarrow \sigma\pi^+)$. To see this, we shall first define a quantity

$$\eta \equiv \frac{\Gamma(D \rightarrow MP \rightarrow P_1 P_2 P)}{\Gamma(D \rightarrow MP)\mathcal{B}(M \rightarrow P_1 P_2)} . \quad (44)$$

As η goes to 1 in the narrow width approximation, the deviation of η from unity gives a measure of violation in the factorization relation (1). We first compute η theoretically; that is, both $\Gamma(D \rightarrow MP \rightarrow P_1 P_2 P)$ and $\Gamma(M \rightarrow P_1 P_2)$ are computed in a model. The ratio η is independent of the form factor for $D \rightarrow M$ transition. The factorization relation Eq. (1) is then replaced by

$$\Gamma(D \rightarrow MP \rightarrow P_1 P_2 P) = \eta \Gamma(D \rightarrow MP)\mathcal{B}(M \rightarrow P_1 P_2) . \quad (45)$$

From the experimental input of $\mathcal{B}(D \rightarrow MP \rightarrow P_1 P_2 P)$, we can then determine $\Gamma(D \rightarrow MP)$.

For $D^+ \rightarrow \sigma\pi^+ \rightarrow \pi^+\pi^+\pi^-$ decays, we find $\eta = 0.55$ for $\Gamma_\sigma = 600$ MeV and $\eta = 0.41$ for $\Gamma_\sigma = 1000$ MeV, where we have taken $\mathcal{B}(\sigma \rightarrow \pi^+\pi^-) = \frac{2}{3}$. This means that the branching fraction $\mathcal{B}(D^+ \rightarrow \sigma\pi^+) = (2.1 \pm 0.2) \times 10^{-3}$ listed in Table I should be enhanced by a factor of $1/\eta = 1.8 \sim 2.4$, depending on the σ width.

$D \rightarrow TP$

We have examined the finite-width effect on the $D \rightarrow TP$ channels listed in Table X. The measured decay widths of various tensor mesons are of order 100 MeV [29].

The singly Cabibbo-suppressed decay $D^+ \rightarrow \overline{K}_2^{*0} K^+ \rightarrow K^+ K^- \pi^+$ is physically allowed due to the width $\Gamma_{K_2^{*0}} = 109 \pm 5$ MeV. From Eq. (38) we obtain

$$\mathcal{B}(D^+ \rightarrow \overline{K}_2^{*0} K^+ \rightarrow K^+ K^- \pi^+) = 4.3 \times 10^{-8} . \quad (46)$$

This is about 3 to 4 orders of magnitude below the experimental result $(1.7_{-0.8}^{+1.2}) \times 10^{-4}$ [29]. However, this measurement seems to be problematic as the branching fraction of the Cabibbo-favored mode $D^+ \rightarrow \overline{K}_2^{*0} \pi^+ \rightarrow K^- \pi^+ \pi^+$ is of the same order of magnitude, $(2.1 \pm 0.4) \times 10^{-4}$.

For the other $D \rightarrow TP$ decays, we compute the ratio η defined in Eq. (44) and find that $\eta \sim 1.0 - 1.2$ for most cases except for $D^0 \rightarrow f_2 \overline{K}^0$ where $\eta = 4.0$. This means that the branching fraction of $D^0 \rightarrow f_2 \overline{K}^0$ extracted in Table III should be reduced by a factor of 4 when the effect of finite width is taken into account⁸.

VII. DISCUSSIONS

A. The covariant light-front model

We have relied heavily on the CLF model to obtain the form factors needed in this work. This relativistic quark model preserves the Lorentz covariance in the light-front framework and has been applied successfully to describe various properties of pseudoscalar and vector mesons [77]. The analysis of the CLF model has been generalized to even-parity, P -wave mesons in [51]. Since

⁸ Our conclusion for finite-width effects on $D \rightarrow TP$ differs from that in Ref. [27].

relativistic effects can manifest in heavy-to-light transitions at large to maximum recoil where the final-state meson becomes highly relativistic, the use of the non-relativistic quark model is probably not suitable here. Therefore, we believe that the CLF approach can provide more accurate behaviors of $B \rightarrow M$ transitions at large recoil. For example, the tensor form factors for $B \rightarrow K_1$ transitions at $q^2 = 0$ derived in the CLF model lead to a prediction for $\mathcal{B}(B \rightarrow K_1(1270)\gamma)$ in much better agreement with experiment than any other models (see Table IV of [56]).

B. Theoretical uncertainties

Thus far we have not given the error bars to the theoretical results of decay rates. Here we discuss the possible sources of uncertainties in this study. In the CLF model we use the decay constants together with the given constituent quark masses to determine a fundamental parameter β in the model for describing the meson wave functions [51]. For the decay constants of scalar mesons we use those obtained in [49] where QCD sum rules are employed and errors are given explicitly. For axial-vector mesons, the sum rule approach gives $f_{a_1} = 238 \pm 10$ MeV [61] while the experimental data of $\tau \rightarrow K_1 \nu$ yield $f_{K_1(1270)} = -(170 \pm 20)$ MeV and $f_{K_1(1400)} = -(139 \pm 43)$ MeV [cf. Eqs. (19) and (20)]. The error bars will be propagated from decay constants to the parameter β , and then finally to the form factors of interest. Therefore, the uncertainty analysis in the CLF model is quite involved and highly nontrivial. We will leave this task to a future publication.

C. Comparison with other works

In this work we have performed the study of the nonleptonic decays of charmed mesons in two approaches: flavor-diagram analysis and naive factorization. In the latter approach, form factors are obtained from the CLF quark model and the weak annihilation diagrams are neglected at the outset. In the literature most of the relevant studies are also based on the factorization approach, differing mainly in the values of the form factors to be used and the treatment of weak annihilation. In Tables IV-VI we have displayed in parentheses the form factors evaluated in the ISGW2 model [52], an improved version of the non-relativistic quark model by Isgur, Scora, Grinstein, and Wise (ISGW) [60]. We see that the ISGW2 model predicts much smaller $D \rightarrow S$ form factors than the CLF model and other models not listed in Table IV. In contrast, the form factors $V_0^{Da_1}$, $V_0^{Db_1}$, k^{Da_2} , $k^{Df_{2q}}$, $k^{Ds f_{2s}}$ and $k^{Ds K_2^*}$ calculated by the ISGW2 model are much larger than the CLF results.

As stressed before, the factorization hypothesis is best tested in the decays in which weak annihilation contributions are absent or suppressed. For the SP modes, the Cabibbo-allowed decays $D^+ \rightarrow \bar{K}_0^{*0} \pi^+$ and $D_s^+ \rightarrow f_0 \pi^+$ belong to this category. It turns out that the former mode is ideal for testing different form factor models. The contribution to $D^+ \rightarrow \bar{K}_0^{*0} \pi^+$ from the color-suppressed tree amplitude C was not considered in [10, 11], presumably due to the smallness of the K_0^* decay constant. Both contributions of color-allowed and color-suppressed amplitudes were taken into account in [9]. However, owing to the destructive interference in $D^+ \rightarrow \bar{K}_0^{*0} \pi^+$ and the smallness of a_2 , the predictions of $\Gamma(D^+ \rightarrow \bar{K}_0^{*0} \pi^+) < \Gamma(D^0 \rightarrow \bar{K}_0^{*-} \pi^+)$ and the large suppression

of $D^0 \rightarrow \bar{K}^{*0}\pi^0$ relative to $D^0 \rightarrow \bar{K}^{*-}\pi^+$ given in [9] are not borne out by experiment. In the CLF model, the relative sign of the factorizable amplitudes $X^{(DS,P)}$ and $X^{(DP,S)}$ are fixed to be negative [see Eq. (29)]. As a consequence, we conclude that, based on the CLF quark model, the interference in the Cabibbo-allowed $D^+ \rightarrow SP$ decays must be constructive, contrary to the case of $D \rightarrow PP, VP$.

Recently, the decays $D \rightarrow K_1\pi$ have been considered in [23, 24] with $D \rightarrow K_1$ form factors evaluated in the framework of the ISGW model and QCD sum rules, respectively. While the predicted rates are similar to ours in most cases, the branching fraction of $D^+ \rightarrow \bar{K}_1^0(1270)\pi^+$ was found to be $(5.85 \pm 0.37)\%$ for $\theta_{K_1} = -37^\circ$ and $(3.18 \pm 0.25)\%$ for $\theta_{K_1} = -58^\circ$ in [24]. All of them are too large compared to the experimental limit 7×10^{-3} [29]. In our study we did include the color-suppressed tree amplitude [see Eq. (33)], which was considered in [23] but neglected in [24]. Since the decay constant of $K_1(1270)$ is negative [Eq. (20)], it is clear from Eq. (33) that the a_2 term contributes destructively to $D^+ \rightarrow \bar{K}_1^0(1270)\pi^+$. Consequently, our prediction $\mathcal{B}(D^+ \rightarrow \bar{K}_1^0(1270)\pi^+) = 4.7 \times 10^{-3}$ (Table IX) is consistent with experiment.⁹

The above few examples indicate that the CLF model takes care of the relative signs of decay constants, magnitudes of form factors, and various hadronic matrix elements correctly.

D. Comparison with B decays

Charmful decays $B \rightarrow D^{**}P$ with $D^{**} = D_0^*, D_1, D_1', D_2^*$ denoting even-parity charmed mesons and charmless decays $B \rightarrow MP$ (M being light even-parity mesons) have been extensively studied both experimentally [29] and theoretically [49, 78–80]. It is instructive to compare the present work with the B decays.

While factorization works well for Cabibbo-allowed $D^+ \rightarrow SP, AP$ decays, predictions are typically about one order of magnitude smaller than experiment for the other decay modes, conceivably due to the negligence of weak annihilation contributions arising from final-state interactions. It is pointed out in [49] that one needs a sizable penguin weak annihilation amplitude in order to account for the data of $B \rightarrow K_0^*\pi$ in the theoretical approach such as QCD factorization.

Taking the cue from the constructive interference in the Cabibbo-allowed decay $D^+ \rightarrow \bar{K}_0^{*0}\pi^+$ and noting that the phase of a_2/a_1 in $B \rightarrow D\pi$ decays lies in the first quadrant, one may be tempted to claim that the $B^- \rightarrow D_0^{*0}\pi^+$ decay should have a rate smaller than the $B^0 \rightarrow D_0^{*-}\pi^+$ decay owing to a destructive interference in the former. This conjecture is supported in both the CLF model and heavy quark symmetry [78]. The experimental observation that the production of broad D^{**} states in charged B decays is more than a factor of five larger than that produced in neutral B decays (see Tables V and VI in [78]) is thus astonishing. This enigma as pointed out several years ago in [78] still remains unresolved.

⁹ $\mathcal{B}(D^+ \rightarrow \bar{K}_1^0(1270)\pi^+)$ is predicted to be 3.8×10^{-3} , 1.52% and 3.21% in [23] for $\theta_{K_1} = 33^\circ, 45^\circ$ and 57° , respectively. However, as noticed in Sec. III.B, the mixing angle θ_{K_1} has to be negative in the convention of QCD sum rules or the ISGW model. Therefore, the mixing angle chosen by [23] does not have a correct sign. Also the relative sign between T and C amplitudes given there is erroneous.

VIII. CONCLUSIONS

In this paper, we have studied the charmed meson decays into final states containing one pseudoscalar meson and one even-parity meson, the data of which are inferred from detailed Dalitz analyses of three-body decays and the finite width effects. The non-perturbative flavor diagram approach and the factorization calculation are undertaken to analyze these decay processes. In the diagrammatic framework, we have extracted the sizes and relative strong phases of various flavor diagrams in a least model-dependent way, based on current experimental measurements of decay rates. In the factorization approach, we have neglected weak annihilation diagrams (E and A) while making our predictions. Besides, we use the form factors evaluated in the covariant light-front model to compute the D decay branching fractions.

In the $D \rightarrow SP$ decays with S being in the non-strange nonet, our fits present several robust features against the 2- and 4-quark pictures. First, as in the PP and VP decays, the weak annihilation diagrams are non-negligible. However, it is a theoretically challenging puzzle that the A amplitude here is the largest one. Secondly, the relative strong phases δ_A and δ_E are preferably around 30° and 160° , respectively. Finally, the data do not prefer either the 2-quark scheme or the 4-quark scheme. This conclusion can be best seen from a comparison between the $D^{0,+} \rightarrow f_0\pi$ and $\sigma\pi$ decays, and also revealed in the $\chi^2_{\min}$ values in Table VIII. We note that due to a paucity of measured modes, the diagrammatic approach is ineffective for the $D \rightarrow K_0^*P$ decays or the $D \rightarrow AP$ decays. Theoretical calculations based on factorization have been applied to both $D \rightarrow SP$ and AP transitions. The factorization hypothesis seems to work pretty well for those Cabibbo-favored modes that involve only the color-allowed tree (T) and color-suppressed tree (C) amplitudes. The factorization calculations for the other decay modes are typically about one order of magnitude smaller than experiment, conceivably due to the negligence of weak annihilation contributions.

Contrary to the Cabibbo-allowed $D^+ \rightarrow PP, VP$ decays where T and C amplitudes interfere destructively, the color-allowed and color-suppressed tree amplitudes in the Cabibbo-allowed decays $D^+ \rightarrow \bar{K}^{*0}\pi^+, \bar{K}_1^0(1400)\pi^+$ contribute constructively. This explains why their branching fractions are large, of $\mathcal{O}(2\%)$.

The $D \rightarrow TP$ measurements poise the biggest problem for theory. Even though the magnitudes and phases of some amplitudes can be extracted from data, as given in Table XI, quite opposite conclusions are reached when different sets of data [Fit (A) versus Fit (B)] are used. This could be caused by the $D^+ \rightarrow \bar{K}_2^{*0}\pi^+$ and $D^+ \rightarrow K_2^{*0}\pi^+$ decays, as explained in the text. Predicted branching fractions based on factorization are at least two orders of magnitude smaller than data, even when the decays are free of weak annihilation contributions. We cannot find possible sources of rate enhancement.

We also examine the finite width effects for decays that are kinematically forbidden if the width of the even-parity meson is not taken into account. We find that the branching fraction of $D^+ \rightarrow \sigma\pi^+$ extracted from three-body decays is enhanced by a factor of 2, whereas $\mathcal{B}(D^0 \rightarrow f_2(1270)\bar{K}^0)$ is reduced by a factor of 4 by finite width effects.

Our study shows that some of the above-mentioned puzzles call for measurements of yet observed decay modes as well as more precise determination in the decay rates. The other puzzles, particularly in the TP modes, demand better understanding of the underlying dynamics.

Acknowledgments

H.-Y. C. wishes to thank the hospitality of the Physics Department, Brookhaven National Laboratory. This research was supported in part by the National Science Council of Taiwan, R. O. C. under Grant Nos. NSC 97-2112-M-008-002-MY3, NSC 97-2112-M-001-004-MY3 and in part by the NCTS.

-
- [1] F. Buccella, M. Lusignoli, G. Miele, A. Pugliese and P. Santorelli, Phys. Rev. D **51**, 3478 (1995) [arXiv:hep-ph/9411286].
 - [2] J. L. Rosner, Phys. Rev. D **60**, 114026 (1999) [arXiv:hep-ph/9905366].
 - [3] M. Gronau, Phys. Rev. Lett. **83**, 4005 (1999) [arXiv:hep-ph/9908237].
 - [4] M. Gronau and J. L. Rosner, Phys. Lett. B **500**, 247 (2001) [arXiv:hep-ph/0010237].
 - [5] C. W. Chiang and J. L. Rosner, Phys. Rev. D **65**, 054007 (2002) [arXiv:hep-ph/0110394].
 - [6] C. W. Chiang, Z. Luo and J. L. Rosner, Phys. Rev. D **67**, 014001 (2003) [arXiv:hep-ph/0209272].
 - [7] M. Ablikim, D. S. Du and M. Z. Yang, High Energy Phys. Nucl. Phys. **27**, 759 (2003) [arXiv:hep-ph/0211413].
 - [8] Z. Ligeti, M. Luke and A. V. Manohar, arXiv:1003.1351 [hep-ph].
 - [9] F. Hussain, A. N. Kamal and S. N. Sinha, Z. Phys. C **36**, 199 (1987).
 - [10] A. C. Katoch and R. C. Verma, Z. Phys. C **62**, 173 (1994).
 - [11] S. Fajfer, Z. Phys. C **68**, 81 (1995).
 - [12] F. Buccella, M. Lusignoli and A. Pugliese, Phys. Lett. B **379**, 249 (1996) [arXiv:hep-ph/9601343].
 - [13] H. Y. Cheng, Phys. Rev. D **67**, 034024 (2003) [arXiv:hep-ph/0212117].
 - [14] B. El-Bennich, O. Leitner, J. P. Dedonder and B. Loiseau, Phys. Rev. D **79**, 076004 (2009) [arXiv:0810.5771 [hep-ph]].
 - [15] D. R. Boito, J. P. Dedonder, B. El-Bennich, O. Leitner and B. Loiseau, Phys. Rev. D **79**, 034020 (2009) [arXiv:0812.3843 [hep-ph]].
 - [16] A. N. Kamal and R. C. Verma, Phys. Rev. D **45**, 982 (1992).
 - [17] X. Y. Pham and X. C. Vu, Phys. Rev. D **46**, 261 (1992).
 - [18] T. N. Pham, Phys. Rev. D **46**, 2976 (1992).
 - [19] A. N. Kamal, Q. P. Xu and A. Czarnecki, Phys. Rev. D **49**, 1330 (1994).
 - [20] A. C. Katoch and R. C. Verma, J. Phys. G **21**, 525 (1995).
 - [21] H. J. Lipkin, Phys. Lett. B **515**, 81 (2001) [arXiv:hep-ph/0011228].
 - [22] H. Y. Cheng, Phys. Rev. D **67**, 094007 (2003) [arXiv:hep-ph/0301198].
 - [23] N. Sharma and R. C. Verma, Phys. Rev. D **75**, 014009 (2007).
 - [24] R. Khosravi, K. Azizi and N. Ghahramany, Phys. Rev. D **79**, 036004 (2009) [arXiv:0812.1352 [hep-ph]].
 - [25] A. C. Katoch and R. C. Verma, Phys. Rev. D **49**, 1645 (1994) [Erratum-ibid. D **55**, 7315 (1997)].

- [26] J. H. Munoz, A. A. Rojas and G. Lopez Castro, Phys. Rev. D **59**, 077504 (1999) [arXiv:hep-ph/9812274].
- [27] H. Y. Cheng, Phys. Rev. D **68**, 014015 (2003) [arXiv:hep-ph/0303195].
- [28] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **74**, 032003 (2006) [arXiv:hep-ex/0605003].
- [29] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [30] E687 Collaboration, P.L. Frabetti *et al.*, Phys. Lett. B **351**, 591 (1995); *ibid.* B **331**, 217 (1994).
- [31] E791 Collaboration, E.M. Aitala *et al.*, Phys. Rev. Lett. **89**, 121801 (2002); *ibid.* **86**, 765 (2001); *ibid.* **86**, 770 (2001).
- [32] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **79**, 032003 (2009) [arXiv:0808.0971 [hep-ex]].
- [33] M. Gourdin, Y. Y. Keum and X. Y. Pham, Phys. Rev. D **53**, 3687 (1996) [arXiv:hep-ph/9506225].
- [34] M. Ablikim *et al.* [BES Collaboration], Phys. Rev. D **72**, 092002 (2005) [arXiv:hep-ex/0508050].
- [35] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **78**, 034023 (2008) [arXiv:0804.2089 [hep-ex]].
- [36] R. E. Mitchell *et al.* [CLEO Collaboration], Phys. Rev. D **79**, 072008 (2009) [arXiv:0903.1301 [hep-ex]].
- [37] C. Amsler and N. A. Törnqvist, Phys. Rept. **389**, 61 (2004).
- [38] F.E. Close and N.A. Törnqvist, J. Phys. G **28**, R249 (2002) [hep-ph/0204205].
- [39] T. Barnes, Phys. Lett. B **165**, 434 (1985).
- [40] D. Black, M. Harada and J. Schechter, Phys. Rev. D **73**, 054017 (2006).
- [41] R.L. Jaffe, Phys. Rev. D **15**, 267 (1977); *ibid.* 281 (1977).
- [42] S. Prelovsek, T. Draper, C. B. Lang, M. Limmer, K. F. Liu, N. Mathur and D. Mohler, PoS **LAT2009**, 103 (2009) [arXiv:0910.2749 [hep-lat]]; S. Prelovsek and D. Mohler, Phys. Rev. D **79**, 014503 (2009) [arXiv:0810.1759 [hep-lat]].
- [43] N. Mathur *et al.*, Phys. Rev. D **76**, 114505 (2007) [arXiv:hep-ph/0607110].
- [44] R. L. Jaffe, Phys. Rept. **409**, 1 (2005) [Nucl. Phys. Proc. Suppl. **142**, 343 (2005)] [arXiv:hep-ph/0409065].
- [45] K. M. Ecklund *et al.* [CLEO Collaboration], Phys. Rev. D **80**, 052009 (2009) [arXiv:0907.3201 [hep-ex]].
- [46] T. M. Aliev and M. Savci, arXiv:hep-ph/0701108.
- [47] L. Maiani, F. Piccinini, A.D. Polosa, and V. Riquer, Phys. Rev. Lett. **93**, 212002 (2004).
- [48] W. Wang and C. D. Lu, arXiv:0910.0613 [hep-ph].
- [49] H.Y. Cheng, C.K. Chua, and K.C. Yang, Phys. Rev. D **73**, 014017 (2006).
- [50] M. Wirbel, B. Stech, and M. Bauer, Z. Phys. C **29**, 637 (1985); M. Bauer, B. Stech, and M. Wirbel, *ibid.* **34**, 103 (1987).
- [51] H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. D **69**, 074025 (2004) [arXiv:hep-ph/0310359].
- [52] D. Scora and N. Isgur, Phys. Rev. D **52**, 2783 (1995).

- [53] M. Suzuki, Phys. Rev. D **47**, 1252 (1993).
- [54] L. Burakovsky and T. Goldman, Phys. Rev. D **56**, 1368 (1997).
- [55] H. Y. Cheng, Phys. Rev. D **67**, 094007 (2003).
- [56] H. Y. Cheng and C. K. Chua, arXiv:0909.4627 [hep-ph].
- [57] H. Y. Cheng and K. C. Yang, Phys. Rev. D **76**, 114020 (2007) [arXiv:0709.0137 [hep-ph]], *ibid.* D **78**, 094001 (2008) [arXiv:0805.0329 [hep-ph]].
- [58] R. H. Li, C. D. Lu and W. Wang, Phys. Rev. D **79**, 034014 (2009) [arXiv:0901.0307 [hep-ph]].
- [59] K. C. Yang, Phys. Rev. D **78**, 034018 (2008) [arXiv:0807.1171 [hep-ph]].
- [60] N. Isgur, D. Scora, B. Grinstein, and M.B. Wise, Phys. Rev. D **39**, 799 (1989).
- [61] K.C. Yang, Nucl. Phys. B **776**, 187 (2007) (arXiv:0705.0692 [hep-ph]).
- [62] D.M. Li, H. Yu, and Q.X. Shen, J. Phys. G **27**, 807 (2001).
- [63] L.L. Chau and H.Y. Cheng, Phys. Rev. D **36**, 137 (1987); Phys. Lett. B **222**, 285 (1989).
- [64] L.L. Chau, Phys. Rep. **95**, 1 (1983).
- [65] L.L. Chau and H.Y. Cheng, Phys. Rev. Lett. **56**, 1655 (1986).
- [66] M. Gronau, O.F. Hernandez, D. London, and J.L. Rosner, Phys. Rev. D **50**, 4529 (1994); Phys. Rev. D **52**, 6374 (1995)
- [67] M. Zhong, Y. L. Wu and W. Y. Wang, Eur. Phys. J. C **32**, 191 (2004).
- [68] Y. L. Wu, M. Zhong and Y. F. Zhou, Eur. Phys. J. C **42**, 391 (2005) [arXiv:hep-ph/0405080].
- [69] B. Bhattacharya and J. L. Rosner, Phys. Rev. D **77**, 114020 (2008) [arXiv:0803.2385 [hep-ph]].
- [70] B. Bhattacharya and J. L. Rosner, Phys. Rev. D **79**, 034016 (2009) [arXiv:0812.3167 [hep-ph]].
- [71] M. Artuso *et al.* [CLEO Collaboration], Phys. Rev. D **77**, 092003 (2008) [arXiv:0802.2664 [hep-ex]].
- [72] B. Bhattacharya and J. L. Rosner, arXiv:0911.2812 [hep-ph].
- [73] D. Besson *et al.* [CLEO Collaboration], Phys. Rev. D **80**, 032005 (2009) [arXiv:0906.2983 [hep-ex]].
- [74] H. Y. Cheng and C. K. Chua, Phys. Rev. D **69**, 094007 (2004) [arXiv:hep-ph/0401141].
- [75] H. Y. Cheng, Phys. Rev. D **67**, 054021 (2003) [arXiv:hep-ph/0212361].
- [76] H. Pilkuhn, *The Interactions of Hadrons*, Amsterdam: North-Holland (1967).
- [77] W. Jaus, Phys. Rev. D **60**, 054026 (1999).
- [78] H. Y. Cheng and C. K. Chua, Phys. Rev. D **74**, 034020 (2006) [arXiv:hep-ph/0605073].
- [79] H. Y. Cheng, C. K. Chua and K. C. Yang, Phys. Rev. D **77**, 014034 (2008) [arXiv:0705.3079 [hep-ph]].
- [80] H. Y. Cheng and K. C. Yang, Phys. Rev. D **76**, 114020 (2007) [arXiv:0709.0137 [hep-ph]].